

Chris Anthony

HOME

NEWS & WEATHER

UNIVERSITY

SCIENCE *General* *My research*

INDIA *General* *My pictures*

MUSIC

FAVOURITES shopping, friends, DIY, entertainment, Southampton etc

BOOKS

TRAVEL & HOLIDAYS

FAMILY

MEDICAL

PHOTOGRAPHY

BIRDS

SEARCH ENGINES

All engines
Google
YouTube
Stumbleupon

MY NET PROVIDER

NOTE 'My files' open with :
ppt Powerpoint
doc Word
xls Excel
pdf Acrobat Reader [download free]
jpg Opens direct
html Opens direct;
For **BIGGER** text
Click Ctl plus +
Normal text click Ctl plus 0

Presentations of some of my Teaching and Research Topics

Go Back to:

[My main Research Page](#) [Science Education page](#) [Methanotroph history](#) [Main Science Page](#) [India Page](#)

There are two sets of material here:

- a) A summary of most of my work on methylotrophs given in a lecture in MAY 2024
- b) The slides taken from my powerpoint presentations as a guest of Prof. Colin Murrell at the University of East Anglia. They include many pictures of relevant scientists. This can be downloaded as a Powerpoint presentation.
- c) Lecture notes and slides for use of students on my Microbial Physiology course in SV University, India. This material can be downloaded as a Word document or as a powerpoint presentation.

[Methylotroph stories: my methylotroph research summarised in about 50 slides PPT](#)

The Methylotrophs and their investigators	PPT	History of Methanotrophy	
Oxidation of methane methanol & formaldehyde	PPT	The oxidation of methylated amines	PP
The cytochromes and electron transport in methylotrophs	PPT	Growth yields of methylotrophs	PF
Carbon assimilation pathways	PPT	Completion of Serine cycle	PF

[Link to: Teaching page on Microbial Physiology \(SV University, India\)](#)

Methylotroph stories; dedicated to research students everywhere (May 2024)

VIDEOS can be speeded up by
downloading an [accelerator](#)

HOME



**This is a power point presentation of a lecture I gave on
20th May to the Microbiology group in Biological Sciences,
University of Southampton**

The lecture summarised my 40 years of work on methylotroph metabolism in 40 minutes.

I have added a few extra slides to make the stories complete.

Methyloleptroph stories

Dedicated to research students everywhere

My CV

Born Watford in 1939, son of a brewery labourer.

1950 – 1957 Watford Grammar School

1957 – 1960 Reading University BSc Microbiology

1960 – 1963 Reading University PhD Thesis: The microbial metabolism of one-carbon compounds

1963 – 1966 Reading University Post Doc: A novel methanol dehydrogenase

1966 - 1967 Sheffield University Post Doc: glutamate fermentation in Clostridia

1967 – 2004 Southampton University: Lecturer etc

2005 – present time: Southampton University: Emeritus Professor

Thank you to my research students (1969 – 2008)

Methanol dehydrogenase	Cytochromes and electron transport	Metabolic pathways	3D structures
Steven Ford	Jim Elliott	Pat Dunstan	Peter James
Dudley Page	Steven Froud	Iain Taylor	Paul Williams
Darren Day	David O'Keefe	John Bolbot	Gyles Cozier
Jonathan Cox	Andrew Cross		Simon Dales
Ian Richardson	Mathew		[John Read]
Alain Avezoux	Beardmore-Gray		[Chenyi]
Karen Amaratunga	Anthony Long		[Leigh Felton]
Tosin Majekodunmi	Claude Chan		
Paul Afolabi	Helen Ashworth		
	Also Bill Keevil (Post-doc); Paul Skipp (Technician)		
Amino acid transport	Electron transport and nitrogen fixation	Neomycin biosynthesis	Electron transport in methylamine metabolism
Jeff Robinson	Andrew Smith	Cedric Pearce	Ashley Lawton
Bob Cook	Nick Juty		Kevin Auton

Methylotroph stories

Methylotrophs are bacteria that grow with compounds with a single carbon atom as their sole source of carbon and energy. 'Ordinary bacteria' never grow methylotrophically.

1. The rise of the methylotrophs. Only 2 species in 1960. Many hundreds now – with enormous genetic and metabolic diversity.
2. Methanol dehydrogenase (**MDH**); the first Quinoprotein.
3. A special electron transport chain for methanol oxidation.
4. A remarkable resurgence of interest in MDH.

My PhD project (1960 – 1963): Isolate a bacterium able to grow on methanol and therefore able to oxidise it. Then find the enzyme responsible.

1. The rise of the methylotrophs.

Always specialised genera; never the usual types of bacteria
Many grow well on 'ordinary' substrates – succinate, lactate etc

Only 2 species in 1960. Only 4 papers in my PhD introduction

Many hundreds of species now – with enormous genetic diversity

A wealth of novel metabolic pathways for conversion of formaldehyde to cell material

Unusual energy transduction


Regular international symposia (from 1970 to present day)

Industrial exploitation. This is because of their metabolic diversity and because their growth substrate, methanol, is inexpensive

The main 'workhorses' in the study of methylotrophs

- *Pseudomonas* AM1 (Peel & Quayle, 1961)
- *Pseudomonas* sp. M27 (Anthony & Zatman, 1964)

Methylobacterium extorquens. A pink facultative methylotroph



Me in 1963



J Rod Quayle




My PhD supervisor and my first research student (Len Zatman and Pat Dunstan, now Goodwin)



Young Howard & Rod Quayle

Howard Dalton, Rod and me

The Biochemistry of Methylootrophs 1982



The Biochemistry of Methylootrophs

C. ANTHONY

*Department of Biochemistry
University of Southampton
England*

Chapters 12

Text pages 350

Index 48 pages

Tables 60

Figures 57

References 1300



ACADEMIC PRESS


A Subsidiary of Harcourt Brace Jovanovich, Publishers

London · New York
Paris · San Diego · San Francisco
São Paulo · Sydney · Tokyo · Toronto

Methylootrophic bacteria
Ribulose biphosphate path
Ribulose monophosphate cy
The Serine pathway
The TCA cycle and growth
on multicarbon compound
Oxidation of methane, metha
formaldehyde and format
Oxidation of methylated ami
Electron transport and
energy transduction
Growth yields and bioenerge
Methylootrophic yeasts
Methanogens and
methanogenesis
Commercial exploitation

A CD version is available and the complete b
is on my website: chris-anthony.co.uk

The ICI 'Pruteen' plant at Billingham, UK 1980 For production of protein supplement for cattle feed



The centre tower is the 1.5 million litre fermenter

It contains about 100,000,000,000,000,000,000 methylotrophs
10% of their soluble protein is Methanol dehydrogenase

2nd Story: Methanol dehydrogenase

An enzyme with a remarkable structure and novel type of prosthetic group - PQQ (the first Quinoprotein). The false claim that PQQ is a new vitamin

Oxidising enzymes.

Most common: alcohol + NAD⁺ → aldehyde + NAD + H⁺

Alternative oxidising enzymes are membrane bound and have FAD or FMN as a bound prosthetic group: succinate + Enz-FAD → fumarate + Enz-FADH₂

Most living things have an NAD-linked alcohol dehydrogenase
BUT none oxidises methanol so my supervisor, Len Zatman, thought there might be an interesting dehydrogenase in bacteria able to oxidise methanol to formaldehyde.

My project: Isolate a bacterium able to grow on methanol and therefore able to oxidise it. Then find the enzyme responsible.

I thought this was a trivial problem but it was not.


My first paper was on a novel methanol dehydrogenase (1964)

My last paper was on the very high (atomic) resolution of this enzyme (2004)

We published 2 papers together. The first described our new methylotroph and the next slide is the second.

Methanol dehydrogenase

Responsible for all bacterial methanol oxidation



A soluble enzyme – now known to be in the periplasm

Dye-linked assay (PMS/PIP); pH optimum 9.2; pI 8.2


Methanol + PMS $\rightarrow$ Formaldehyde + PMSH₂
PMSH₂ is oxidised by blue PIP which loses its colour

Activator is ammonia (not NH₄⁺ ion) (this is unique)

Wide substrate specificity

Unusual absorption spectrum – not FAD or FMN **so novel prosthetic group??**

High proportion of tryptophans



MDH assay
with PMS/indophenol

Isolation of the prosthetic group

The prosthetic group is the part of the enzyme that is not the amino acid chain.

It is FAD or FMN in flavoproteins, or haem in cytochromes

Grow huge amounts of bacteria on methanol. Harvest and break (ultrasound)
Purify the enzyme from the soluble fraction (Ion exchange, Sephadex etc)

Precipitate the protein and release the fluorescent prosthetic group by acid treatment

Purify the prosthetic group (ion exchange etc). After 3 months I had a few red grains.

Characterised by its excitation and emission fluorescence. A beautiful green colour.


The only known compound with a maximum excitation at 360nm with this maximum occurring at low pH. It is tightly bound to the enzyme.

SO a completely new compound. Thirteen years later its structure was determined by X-ray crystallography at Cambridge; then confirmed by us a further 15 years later with the X-ray structure of the whole enzyme.


A new type of enzyme prosthetic group

Note: all other dehydrogenases use NAD(P) as hydrogen acceptor or they are membrane flavoproteins that use FMN or FAD as prosthetic group

The Microbial Oxidation of Methanol
 THE PROSTHETIC GROUP OF THE ALCOHOL DEHYDROGENASE OF
PSEUDOMONAS SP. M27:
 A NEW OXIDOREDUCTASE PROSTHETIC GROUP
 BY C. ANTHONY* AND L. J. ZATMAN
Department of Microbiology, University of Reading
 (Received 30 December 1966)



Spectrum of MDH



Spectrum of prosthetic group

PQQ and other quinoproteins The novel prosthetic group of MDH was later in other dehydrogenases and Hans Duine and Hans Frank called these Quinoproteins. These include the alcohol dehydrogenase of acetic acid bacteria and the typical glucose dehydrogenase in *Pseudomonas*, *E. coli*.



Hauge described GDH and PQQ very early on




Hans Duine and Hans Frank
'invented' quinoproteins

Friends from Yamaguchi, Japan:
Matsushita, Uragami (?), Ameyama, Adachi.
Found many new quinoproteins especially in
Acetic acid bacteria

**Note: No PQQ-containing quinoproteins
have been described in anything other
than bacteria**

PQQ and Quinoproteins

Pyrroloquinoline Quinone PQQ)



First isolated and purified
by Anthony & Zatman
in 1967 from bacterial
methanol dehydrogenase

Structure determined by
Kennard's group in 1979

Remains tightly bound
throughout reaction cycle

PQQ-containing Quinoproteins

All bacterial dehydrogenases

Usually responsible for 1st step in
the oxidation of growth substrate

Mainly alcohols and aldoses

eg ethanol oxidation to acetic acid
in acetic acid bacteria

Reaction occurs in the periplasm

Electron acceptors are
cytochromes, blue copper proteins
or ubiquinone

May be soluble or membrane-bound

May also have contain haem

**None has been described in
eukaryotic organisms**

From 1988 onwards there have been International Symposia on PQQ and Quinoproteins The 5th was in Southampton in 2002




3rd International Symposium on Vitamin B₆, PQQ, Carbonyl Catalysis and Quinoproteins



University of Southampton, UK

April 14th-19th 2002



Comprising the
11th International Symposium on Vitamin B₆ and Carbonyl Catalysis
and the 5th International Symposium on PQQ and Quinoproteins



Boldrewood conference Centre

Programme book and Abstracts

The following slide was included to show what was lost when Boldrewood was demolished. This was my room, complete with coffee machine and comfort. Murali with back showing was a 4th year medical student solving a problem with his project and the other is a research student friend Leigh Felton. Staff rooms were always available for students to see members of staff.



My office with Murali and Leigh


The following section describes our work with MDH which was done some years later in collaboration with Meenakshi Ghosh who came to Oxford from Calcutta to learn how to do X-ray crystallography of proteins. She had previously had experience only with inorganic crystals. She needed a project so I provided it.

The X-ray structure of MDH 1992-2004

Southampton:
Chris Anthony
Alain Avezoux

Oxford:
Meenakshi Ghosh,
Colin Blake


MDH is a dimer
Large subunit is 60k
Small subunit is 9k



Meenakshi
Ghosh

A wonderful collaboration; Meenakshi knew no biochemistry and i knew no crystallography and its interpretation

The β -propeller structure of MDH



We made the mistake of relying on an 'expert' to assay our enzyme for metal ion content and he found none. After our Japanese friends showed that their Ethanol dehydrogenase contained calcium we found it in MDH. Our work with a mutant lacking calcium enabled us to reconstitute it with strontium or barium.

The calcium ion in MDH


1965-1989 'MDH contains no metal ion' [Anthony & Zatman]

1986 MDH from mutants (*mxmA*, *K* & *L*) was inactive and had unusual spectrum [Nunn & Lidstrom]

1990 MDH contains Calcium (or strontium) [Adachi et al.]

1992 Other MDHs contain calcium.
MDH from *mxmA*, *K* & *L* lacked Calcium;
incubation with Ca produced active enzyme
& normal spectrum [Ian Richardson]


1992 MDH isolated from marine *Methylophaga Marina* is red and inactive;
incubation with Ca produced
normal green active enzyme [Claude Chan]




The calcium ion in MDH

1994 Kinetics of MDH containing strontium *[Harris & Victor Davidson]*

1996 Incubation with Barium produced MDH which was more active than wild-type MDH [Matt Goodwin, Alain Avezoux & Simon Dales]










Tryptophan docking system

High proportion of tryptophans;
19 per α subunit

The β -sheet 'propeller blades' are held together by 'tryptophan-docking motifs'



The following slide shows the consensus sequence involved in the docking motifs which demonstrated to Kashahara and Kato that they had a propeller structure which they wrongly interpreted as evidence for a quinhop


The consensus sequence in the tryptophan-docking motifs

Position	1	2	3	4	5	-	6	7	8	9	10	11
Motif	C	C	C				D	D	D	D	D	D
W1	Ala 77	Leu	Gly	Leu	Asp	Asp	Pro	Gly	Thr	Ile	Leu	Trp 88
W2	Ala135	Leu	Asn	Ala	Glu	-	Thr	Gly	Glu	Thr	Val	Trp145
W3	Ala 186	Tyr	Asp	Val	Lys	-	Thr191-Gly192		Glu	Gln	Val	Trp 196
W4	Gly 282	Arg	Asp	Ala	Asp	-	Thr	Gly	Glu	Ala	Lys	Phe 292
W5	Thr 337	Leu	Asp	Arg	Thr	-	Asp	Gly	Ala	Glu	Val	Ser 347
W6	Ala 457	Tyr	Asn	Ala	Ile	-	Thr	Gly	Asp	Tyr	Lys	Trp 467
W7	Ala 489	Arg	Asp	Ser	Asp	-	Thr	Gly	Asp	Leu	Leu	Trp 508
W8	Val 577	Phe	Ser	Leu	Asp 581	-	Gln 39	Leu	Arg	Pro	Ala	Trp 44
MDH Consensus	Ala	X	Asp/ Asn	X	X	-	Thr	Gly	Asp/ Glu	X	X	Trp
GDH/ADH Consensus	Ala	X	Asp/ Asn	X	X	-	Thr	Gly	Lys	X	X	Trp

Propeller blades W1-W8

For example: **Trp145** of strand D in W2 is stacked between **Ala135** in strand C in W2 and **Thr-Gly** at position **6-7** in W3


The Direct hydride transfer mechanism Anthony et al (1993--)



The mechanism involves an initial proton abstraction (by Asp 303) and a hydride transfer from the methyl group of methanol. The quinol then passes its electrons one at a time to the specific cytochrome electron acceptor. After a single electron has been transferred the free radical PQQH is stable, and MDH is usually isolated with PQQ in this form (Shown by Frank and Duine)

The unique disulphide ring structure in the MDH active site


Two distant Cysteine residues in proteins can be oxidised to form a disulphide bridge that is used to stabilise a particular conformation





Ramachandran stated that 2 adjacent cysteines would never form a bridge. BUT if they did then they would form an 8-membered ring structure with a 'strained' peptide bond.

**None has ever been seen except in the active site of MDH.
It is very easily reduced, so breaking the bond and inactivating the enzyme.**

The active site of MDH



Novel disulphide ring structure from adjacent cysteines




The path of electrons and protons during the oxidative part of the reaction mechanism of MDH

Electrons must pass from reduced PQQ (PQQH₂) to a unique type of cytochrome - cytochrome c_L

Note involvement of the disulphide ring

Protons must pass from PQQH₂ by way of its hydroxyls, and the hydrogen-bonding network, to the water (W904) which is accessible to the periplasm

The pathways are based on those proposed for the quinohaemoprotein alcohol dehydrogenase by Oubrie, Dijkstra *et al*, and Chen, Matsushita, Toyama, Scott Mathews *et al*.



False claim that PQQ is vitamin

Most enzyme cofactors cannot be made in humans so have to be provided as their precursors – vitamins. Eg: NAD (Niacin), FAD (riboflavin)

If mammals do have quinoprotein enzymes then PQQ is almost certainly a vitamin.

In 2003 Kashara and Kato published a paper in Nature claiming to have found a gene coding for a quinoprotein and announcing that this meant that PQQ is “The first new vitamin for 55 years”.

This was welcomed with a great fanfare

Their argument was so obviously false so I ignored it.

BUT two years later Mitsubishi announced that they were investing 4 million dollars in producing this new vitamin. So I contacted them and published a paper in Nature explaining why it is not. But Mitsubishi went ahead and now it is on the market as a growth nutrient!

Premature announcement that PQQ is a vitamin for mammals

brief communications

Nutritional biochemistry

A new redox-cofactor vitamin for mammals

Takaoki Kasahara, Tadafumi Kato
*Laboratory for Molecular Dynamics of
 Mental Disorders, Brain Science Institute,
 RIKEN, Japan*



NATURE | VOL 422 | 24 APRIL 2003 | www.nature.com/nature

Scientists Add New Member to Vitamin B Family

---Sarah Graham, Scientific American, April 24, 2003.

A new page may need to be added to the vitamin B family album.

According to a report published today in the journal Nature, researchers have determined that the compound known as pyrroloquinoline quinone (PQQ) should be classified as a vitamin in the same class as vitamins E and B3. Scientists first isolated PQQ from bacteria in 1979.



MITSUBISHI GAS CHEMICAL COMPANY, INC.

Biological Products

September 24, 2003

Mitsubishi Gas Chemical Company Inc. speeds the development of
 application of coenzyme PQQ-vitamin.

IN THE NEWS

Japanese Discover First New Vitamin in 55 Years

April 24, 2003
 By REUTERS

**Mind your PQQ - properties of pyrroloquinoline quinine,
 nutrient - vitamins and supplements in a bodybuilder's di**

The argument by Kasahara and Kato

Search engines 'identified' PQQ-binding sites in a protein in mice [predicted from a gene sequence]

Our argument against

We cannot identify PQQ-binding sites in a database

In fact, the search engine found 7 'propeller blades' that typically occur in PQQ-containing quinoproteins

The 'propeller blades' – wrongly identified by the search engine as 'PQQ-binding sites' are not directly involved in binding PQQ; they are purely structural



All trees are green

All PQQ enzymes have propeller structure

My coat is green

My protein has a propeller structure

SO: My coat is a tree

SO: My protein is a PQQ enzyme

Therefore PQQ is a vitamin

brief communications

Nutritional biochemistry

A new redox-cofactor vitamin for mammals

Premature announcement that PQQ is a vitamin for mammals

Leigh M. Felton and Chris Anthony:
Nature; Communication Arising

The only evidence that the proposed dehydrogenase is PQQ-dependent is that the Smart and Pfam databases 'identified' '7 PQQ-binding sites' in the C-terminal region.


In fact, the 'sites' are sequences predicted to form a series of β -sheet 'blades' in a 'propeller fold' as seen in PQQ-dependent dehydrogenases.

This fold has nothing to do with PQQ binding and is found in many other proteins.

The proposed enzyme has no sequence similarity to any PQQ-dependent dehydrogenase.

There is still no evidence that PQQ
Is a cofactor for a eukaryotic enzyme
Or that it is a vitamin

Thank you,
Goodbye and Good luck





REGARDED AS A SUPERCHARGED COQ10, PQQ IS AN ALL NATURAL NUTRIENT THAT PLAYS AN IMPORTANT ROLE IN ENERGY PRODUCTION

Nutritional information

Serving Size: 1 capsule
Servings Per Container: 60

Ingredient	Amount Per Serving	%DV*
pyrroloquinoline quinone (as PQQ disodium salt)	30 mg	**

*Percent Daily Values (DV) are based on a diet of other people's secrets. **No DV established.

INGREDIENTS:
Pyrroloquinoline Quinone (Fuling agent), vegetable magnesium stearate (for vegetable binding), hydroxypropyl methylcellulose (Coating).

DIRECTIONS FOR USE:
Take one capsule daily, preferably with a meal. Do not exceed stated dose.

Best before End: 05/2022
Lot: PQQ097
516603819533400

Triple Strength PQQ (Pyrroloquinoline Quinone) 30 x 60 Veggie Capsules

★★★★★ 11 reviews ⌚ Delivery in 1-2 days 🚚 Free Delivery

Get a daily dose of this all new super potent anti-oxidant, regarded by many as Supercharged CoQ10. Our super strength PQQ has a potent 30mg of the finest-quality Pyrroloquinoline Quinone, triple the strength of most PQQ supplements.



Available options (please select one)

1 Bottle - £27.99 Per Bottle	2 Bottles £22.39 Per Bottle
4 Bottles £20.99 Per Bottle	6 Bottles £18.19 Per Bottle

1 £27.99

3rd Story. Before we knew the structure of MDH we started working on how this enzyme with a novel prosthetic group interacts with an electron transport chain so that the oxidation of methanol will be coupled to ATP production.

3rd Story: A special electron transport chain for methanol oxidation involves a novel type of cytochrome c

Methylobacterium has a 'normal' system for NADH and succinate oxidation

Inhibitors of NADH and succinate oxidation **did not** inhibit methanol oxidation

Inhibitors of the cytochrome oxidase **did** inhibit methanol oxidation


Addition of lactate (giving NADH) or succinate reduced cytochromes b, c and oxidase

BUT: Addition of methanol only reduced cytochromes c and oxidase

Methylobacterium and all methylotrophs have an extra cytochrome c called cytochrome c_L . This is the specific electron acceptor for MDH. They react in the periplasm.

A student David O'Keefe boldly went against my advice, staying up most of the night and found it after months of my failure.

Electron transport during methanol oxidation



During oxidation of MeOH 2 protons are released into the periplasm and consumed inside, setting up a protonmotive force supporting production of about 0.6 ATP/MeC


1975 - 1985 David O'Keefe; Sandy Cross; Mathew Beardmore-Gray; Claude Chan; Jon Cox; Darren Day; Steve Froud; Andy Elliott; Bill Keevil; Sasha Netrusov; Pat Goodwin

NOTE: The ATP yield from methanol is very low but this does not matter because almost every atom of carbon in these bacteria comes by way of this oxidation process so there is ATP to spare. Cell yields are not limited (as in other bacteria) by ATP but by availability of NADH. Matt Beardmore-Gray calculated that the large amount of cytochrome and MDH in these bacteria is sufficient to cover the surface in a mono-molecular layer.

The solution of the unusual MDH structure helped us get an X-ray crystallography unit set up in the Department,

with the appointment of John Cooper and Steve Wood, and they helped solve the structure of the two methylotroph cytochromes.

Structure of cytochrome c_H



M. extorquens
cytochrome c_H

Mammalian
cytochrome c


Bacterial
cytochrome c_2

Remarkably, cytochrome c_H is unique for a bacterial cytochrome in being similar in structure to mammalian cytochrome c

Its internal structure 'disproves' a key proposal for cytochrome function: All other cytochromes c have a tyrosine bonded to water in the haem cleft: Cytochrome c_H has tryptophan and no water. So the water has no function.

Paul Williams and Southampton X-ray structure group

NOTE: the unique calcium ion and the exposed haem propionates



4. A resurgence of interest in MDH

In 1986 Mary Lidstrom and David Nunn published the methanol operon.


David came to work in my lab and he sequenced the genes for MDH and cytochrome c_L

A very remarkable observation

Mary Lidstrom showed that all methylotrophs have two methanol oxidation operons.

The 2nd (Mox2) is never expressed. Mutations in Mox2 do not affect growth on methanol.

Expression of the mxa operon



The genes: Nunn, Lidstrom, Amaratunga, Anderson, Anthony, Goodwin, Morris, O'Connor

NOTE: the immense amount of difficult work on the genetics of our *Methylobacterium* by Karen Amaratunga and Simon Dales led them to provide important evidence for all our ideas about structure, mechanism and electron transport by a programme of site-directed mutagenesis.

A novel methanotroph (Methanotrophs grow on methane which is oxidised to methanol which is then oxidised by MDH)

In 2017 a novel methanotroph (*Methylacidiphilum fumariolicu*) was isolated from acidic volcanic mud. It grows at high temperature and low pH.

It requires a rare earth element for growth (eg Lanthanum, Cerium, Neodymium).

It only uses the 2nd methanol oxidation operon to produce MDH which uses one of the lanthanides instead of calcium at the active site. As a result during the last few years I have weekly reports on ResearchGate of my 1964 paper being cited!

A final strange thing

In 2018 Colin Murrell's group at UEA described a novel methylotroph isolated from seawater using a methanol medium containing lanthanum salts.

"We propose the name *Marinibacterium anthonyii* (in honour of the British microbiologist Professor Christopher Anthony)". Note: I was Colin Murrell's tutor here and he played second violin in my string quartet.

Marinibacterium anthonyii is the only methylotroph that does **not have 'my' MDH; it only has the 2nd methanol oxidation operon.**

Our last publication on MDH 40 years after my first

Acta Crystallographica Section D
**Biological
 Crystallography**
 ISSN 0907-4449

**P. A. Williams, L. Coates,
 F. Mohammed, R. Gill,
 P. T. Erskine, A. Coker,
 S. P. Wood, C. Anthony and
 J. B. Cooper***


School of Biological Sciences, University of
 Southampton, Bassett Crescent East,
 Southampton SO16 7PX, England

The atomic resolution structure of methanol dehydrogenase from *Methylobacterium extorquens*


The crystal structure of methanol dehydrogenase (MDH) from *Methylobacterium extorquens* has been refined without stereochemical restraints at a resolution of 1.2 Å. The high-resolution data have defined the conformation of the tricyclic pyrroloquinoline quinone (PQQ) cofactor ring as entirely planar. The detailed definition of the active-site geometry has

Received 1 September 2004
 Accepted 23 October 2004

PDB Reference: methanol
 dehydrogenase, 1w6s,
 r1w6ssf.



Acknowledgements




Len Zatman & Pat Goodwin




Kevin Auton



Claude Chan



Darren Day



Matt
Beardmore-Gray



H. Dalton



Paul Skipp



Karen
Amaratunga



Sasha
Netrusov





Hiro
Toyama

Acknowledgements



A few more pictures of methylotroph friends



More friends




The oxidative metabolism of methylotrophic bacteria

These slides were to support (very) informal seminars at UEA as guest of Professor Colin Murrell.


There are separate presentation on cytochromes etc

Almost all the enzymes for the initial oxidation steps to formaldehyde are unique to methylotrophs. These unique enzymes are linked to unusual energy-yielding electron transport chains.



- a) Hydroxylation of methane to methanol [Dalton, Murrell etc]
- b) Oxidation of methanol to formaldehyde [Main topic of this seminar]
- c) Oxidation of methylated amines to formaldehyde [Separate seminar]
- d) Oxidation of formaldehyde to formate [Next slide]
- e) Oxidation of formate to carbon dioxide (typical NAD-linked enzyme)
- f) Electron transport chains for energy production

Hydroxylation of methane to methanol



Catalysed by Methane mono-oxygenase (MMO) (Two types)

Soluble (sMMO)


Membrane bounds (particulate or pMMO)

Mainly solved by Howard Dalton; genetics (etc) by Colin Murrell at University of Warwick

More than half oxygen is used by initial hydroxylation in a reaction that requires input of energy (as NADH)

Important people in study of methane oxidation

The late Sir Howard Dalton FRS. Professor Colin Murrell, Mary Lidstrom, Ed Bellion



My second fiddle (juvenile form of Murrell), when I was his tutor at University of Southampton



Nearly adult form

Oxidation of formaldehyde by methylotrophs

Dye-linked dehydrogenases; so possibly flavoproteins or quinoproteins
 NAD-linked dehydrogenases. Absolutely essential during growth on methane with NADH as donor to the hydroxylase
 Methanol dehydrogenase. Good substrate but not likely to be relevant.
 Complex route involving methanopterin and folates [next slide] (NAD-linked)
 Cyclic route in bacteria with RuMP pathway

Note: There must always be routes for removal of toxic formaldehyde. And many aldehyde dehydrogenases are non-specific





Fig. 25. The cyclic route for oxidation of formaldehyde. This is probably the main route for oxidation of formaldehyde in *Pseudomonas C*, *Methylophilus methylotrophus* and *Pseudomonas oleovorans*. This cyclic pathway is discussed further in Chapter 3.

Formaldehyde oxidation in *Methylobacterium extorquens*



Methanopterin was previously only known as carrier in methanogens. This work is by Lidstrom and colleagues. Much is based on genetic/mutant studies.



Last page of my PhD Thesis (submitted 50 years ago; May 1963) suggesting folate involvement in formaldehyde oxidation

Fig.40.

Hypothetical pathways for the oxidation and assimilation of methanol



The oxidation of methanol in Gram negative methylotrophs

This was my PhD project with Len Zatman (1960-1963). He thought the enzyme for methanol oxidation would be unusual as no NAD-linked alcohol dehydrogenase is able to oxidise methanol.

I started by isolating a soil methylotroph, *Pseudomonas* sp M27. It was the same as Quayle's *Pseudomonas* AM1 (published at end of my first year); now called *Methylobacterium extorquens* AM1

Whole cells oxidised methanol and a very wide range of primary alcohols

Inhibited by p-nitro- phenylhydrazine. (50% inhibition by 1.0 μ M). *This has never been explained or investigated further.*

Inhibited by EDTA. Not due to chelation of metal ions. Unique to alcohol oxidation. Later shown to inhibit competitively the binding MDH to its cytochrome electron acceptor

Extracts did not catalyse reduction by methanol of NAD but it was possible to get reduction of a blue indophenol dye in the presence of an intermediate phenazine methosulphate using whole cells in anaerobic conditions.

We then found that this assay could be used with sonic extracts: this is similar to flavoprotein dehydrogenases like succinate dehydrogenase.

The enzyme is **Methanol dehydrogenase (MDH)**.

Problems with the dye linked system for methanol dehydrogenase

The rate was not proportional to protein concentration.

No dye reduction could be seen with lower concentrations of extract.

When water in the assay was replaced with salts growth medium the rate became proportional to protein concentration.

The active constituent was ammonium chloride NH_4Cl .

The activator species is Ammonia base NH_3 not the ammonium ion NH_4^+

This is unique. In some enzymes ammonium ions replace potassium ions. But with MDH ammonium chloride could only be replaced with methylammonium chloride CH_3NH_3^+

At limiting concentrations of activator 10 times more CH_3NH_3^+ was required than with NH_4^+ . This is because its pK value is 10 times higher than that of ammonium chloride.

AND 10 times more ammonium chloride was required at pH 8, when the proportion of NH_3 was one tenth of the proportion at pH 9.



Methanol dehydrogenase

Responsible for all bacterial methanol oxidation

The Microbial Oxidation of Methanol

2. THE METHANOL-OXIDIZING ENZYME OF *PSEUDOMONAS* SP. M 27

BY C. ANTHONY AND L. J. ZATMAN
Department of Microbiology, University of Reading

(Received 21 November 1963)

Dye-linked assay (PMS/PIP); pH optimum 9.2; pI 8.2

Activator- ammonia (not NH_4^+ ion)


Wide substrate specificity

Unusual absorption spectrum – novel prosthetic group??

High proportion of tryptophans

A homodimer (two 60 kDa subunits); later shown to have two 66kDa subunits and two 8kDa subunits

No metal ion. [Later shown to have Calcium ion]



MDH assay
with PMS/indophenol

A new type of enzyme prosthetic group



Note: all other dehydrogenases use NAD(P) as hydrogen acceptor or they are membrane flavoproteins that use FMN or FAD as prosthetic group

The Microbial Oxidation of Methanol

THE PROSTHETIC GROUP OF THE ALCOHOL DEHYDROGENASE OF
PSEUDOMONAS SP. M27:
A NEW OXIDOREDUCTASE PROSTHETIC GROUP


BY C. ANTHONY* AND L. J. ZATMAN
Department of Microbiology, University of Reading

(Received 30 December 1966)



Spectrum of MDH

PQQ 1979 Kennard's lab




Methanol Dehydrogenase [MDH] 1995

New prosthetic group is PQQ

Pyrrolo-Quinoline Quinone

MDH is now known as a Quinoprotein

Pyrroloquinoline Quinone (PQQ)



First isolated and purified
by Anthony & Zatman
in 1967 from bacterial
methanol dehydrogenase

Structure determined by
Kennard's group in 1979

Quinoproteins are enzymes
with an *ortho*-quinone prosthetic
group

They are equivalent to
flavoproteins which has riboflavin
derivatives FMN and FAD as
prosthetic group

PQQ and other quinoproteins The novel prosthetic group of MDH was later in other dehydrogenases and Hans Duine and Hans Frank called these Quinoproteins. These include the alcohol dehydrogenase of acetic acid bacteria and the typical glucose dehydrogenase in *Pseudomonas*, *E. coli*.



Hauge described GDH and PQQ very early on



Friends from Yamaguchi, Japan:
Matsushita, Uragami (?), Ameyama,
Adachi.
Found many new quinoproteins especially
in Acetic acid bacteria




Hans Duine and Hans Frank
'invented' quinoproteins

**Note: No PQQ-containing quinoproteins
have been described in anything other
than bacteria**





Substrate specificity of methanol dehydrogenase



Substrates oxidised




Only primary alcohols




Substrates not oxidised



Alcohols with C2 substituent
are not oxidised




Surprisingly, propane 1,2-diol is oxidised
But only in crude preparations



Modifier protein for methanol dehydrogenase

This was found by accident when investigating growth on propane 1,2-diol

This is oxidised by MDH but **only in crude extracts**;
it does not obey the substrate rules for MDH




The periplasmic Modifier protein **increases affinity for propanediol**

Surprisingly, it is also present in obligate methylotrophs that do not grow on propanediol

M-protein decreases the $V_{\max}$,
and **decreases affinity, for formaldehyde**

This prevents the wasteful oxidation of methanol through formaldehyde to formate



1980-1990: John Bolbot; Steve Ford; Dudley Page; Tony Long

Energy production from methanol dehydrogenase

For energy production the MDH must be linked to the electron transport chain

The cytochromes of *M. extorquens* AM1

Membranes have typical b-type cytochromes and a cytochrome aa_3 able to oxidise typical soluble mitochondrial cytochrome c.

The Soluble fraction has 2 cytochromes c; one is small typical high pl (cytochrome c_H) which is oxidised by membrane fractions. And a much larger one with low pl (cytochrome c_L) which is not oxidised by the membrane oxidase.

Cytochrome c_L is a unique cytochrome. It is the electron acceptor for MDH. It is then able to pass the electrons on to cytochrome c_H which is oxidised by the membrane oxidase. [There is a separate presentation on these cytochromes]


Proton translocation during methanol oxidation

Measured by typical Peter Mitchell method.

The ratio of protons pumped with pyruvate (NADH), succinate and methanol are 3:2:1.


Conclusion: methanol is oxidised by a unique electron transport chain that is similar to those for oxidation of inorganic ions by chemolithotrophs, where ubiquinone and the cytochrome bc_1 complex are not involved.

Electron transport during methanol oxidation



During oxidation of MeOH 2 protons are 'translocated' from inside to outside
Setting up a protonmotive force supporting production of about 0.6 ATP/MeOH

1975 - 1985 David O'Keefe; Sandy Cross; Mathew Beardmore-Gray; Claude Chan;
Jon Cox; Darren Day; Steve Froud; Andy Elliott; Bill Keevil; Sasha Netrusov; Pat Goodwin



The second subunit of MDH

1965 - 1989 MDH is a homodimer (2 x 60 kDa); 'confirmed' by others for > 20 enzymes

1989 Jim Elliott insisted that his 8.5 kDa 'contaminant' must be a subunit



1986: The key papers starting genetics of the methanol oxidation system


Nunn & Lidstrom (1986a,b) *J. Bacteriol.* **166**

1989 We (David Nunn) sequenced the gene for cytochrome c_L (*mxhG*); and showed that the downstream gene (*mxhI*) in the methanol oxidation operon coded for the 'contaminant' which thus became the 8.5kDa β subunit



1989 Darren Day confirmed that MDH is an $\alpha_2\beta_2$ tetramer

Expression of the mxa operon



The genes: Nunn, Lidstrom, Amaratunga, Anderson, Anthony, Goodwin, Morris, O'Connor

The calcium ion in MDH

1965-1989 'MDH contains no metal ion' [Anthony & Zatman]


1986 MDH from mutants (*mxmA*, *K* & *L*) was inactive and had unusual spectrum [Nunn & Lidstrom]

1990 MDH contains Calcium (or strontium) [Adachi et al.]

1992 Other MDHs contain calcium.

MDH from *mxmA*, *K* & *L* lacked Calcium;
incubation with Ca produced active enzyme
& normal spectrum [Ian Richardson]

1992 MDH isolated from marine *Methylophaga Marina* is red and inactive;
incubation with Ca produced
normal green active enzyme [Claude Chan]



The calcium ion in MDH

1994 Kinetics of MDH containing strontium [*Harris & Victor Davidson*]

1996 Incubation with Barium produced MDH which was more active than wild-type MDH [Matt Goodwin, Alain Avezoux & Simon Dales]



The Mechanism of MDH

1979: X-ray structure of PQQ; *Salisbury, Forrest, Cruse, Kennard*

1978 – 1991: Main work on the mechanism; *Hans Frank and Hans Duine*

Ping-pong reaction with PQQ quinol produced from quinone

Reaction is catalysed by an active site base;

Addition/elimination reaction with a covalently attached methanol




The X-ray structure of MDH (1992 – 2004)



Meenakshi Ghosh, Chris Anthony; Colin Blake, Alain Avezoux, Matt Goodwin
Scott Mathews, Z-X Xia, Victor Davidson, Scott White et al.

The structure of MDH of *M. extorquens*




Meenakshi Ghosh, Chris Anthony; Colin Blake, Alain Avezoux, Matt Goodwin

Scott Mathews, Z-X Xia, Victor Davidson, Scott White et al.

1992 - 2003






The β -propeller structure of MDH




Tryptophan docking system

High proportion of tryptophans;
19 per α subunit



The β -sheet 'propeller blades' are held together by 'tryptophan-docking motifs'



The active site of MDH



Novel disulphide ring structure from adjacent cysteines



The path of electrons and protons during the oxidative part of the reaction mechanism of MDH


Electrons must pass from reduced PQQ (PQQH₂) to cytochrome c_L

Cytochrome c_L

Note involvement of the disulphide ring

Protons must pass from PQQH₂ by way of its hydroxyls, and the hydrogen-bonding network, to the water (W904) which is accessible to the periplasm

The pathways are based on those proposed for the quinohaemoprotein alcohol dehydrogenase by Oubrie, Dijkstra *et al*, and Chen, Matsushita, Toyama, Scott Mathews *et al*.



Movie of MDH Made by [Stuart Findlow](#), University of Southampton, from the X-ray diffraction coordinates of MDH. If you cannot movie when you click on arrow then [Go to YouTube clip](#)


There is a separate set of slides for the special cytochromes of methylotrophs

[Top of Oxidation](#)

[Top of Summaries](#) [Return to History](#)

[The oxidation of methylated amines in methylotrophs](#) [PPT](#) [Top of Summaries](#)

Oxidation of methylated amines




Tetramethylammonium Trimethylamine Dimethylamine Methylamine Ammonia Formaldehyde

Metabolism involves demethylation which always occurs at the same time as an oxidation. One methyl group is removed at a time, the C1 unit being removed as formaldehyde.

Methylated amines in the marine environment: Quote from [Yin Chen](#),¹ [Nisha A. Patel](#), [Andrew Crombie](#), [James H. Scrivens](#), and [J. Colin Murrell](#)¹

Methylated amine compounds, including TMA, are ubiquitous in the environment—for example, as end products of protein putrefaction (18). In the marine environment, methylated amines are released as a result of degradation of quaternary amine osmoregulators, such as glycine betaine, which are used by marine organisms to counteract water stress (19–21). Once released into the environment, methylated amines can be used by microorganisms as a C or N source. In fact, in the oceans, methylated amines represent a significant pool of C and N, and standing concentrations up to hundreds of nanomolar and micromolar have been reported in the water column (22, 23) and sediment pore water (24, 25), respectively. In addition to being involved in biogeochemical cycles of C and N, recent studies also suggest that methylated amines have the potential to affect global climate, being precursors of aerosol formation in the upper atmosphere (26–28




(a) Tetramethylammonium monooxygenase; (b) trimethylamine dehydrogenase (in obligate methylotrophs); (c) trimethylamine monooxygenase (in facultative methylotrophs); (d) trimethylamine TV-oxide demethylase (aldolase); (e) dimethylamine monooxygenase (all methylotrophs); (f) dimethylamine dehydrogenase (anaerobic *Hyphomicrobia*); (g) amine oxidase (in *Arthrobacter* and methazotrophic yeasts); (h) methylamine dehydrogenase; (i) N-methylglutamate synthase; (j) and (k) N-methylglutamate dehydrogenases.

Bacterial oxidation of methylated amines

Much of the earlier work on this was done by Bob Eady in the lab of Peter Large in Hull in *Pseudomonas* AM1 (now *Methylobacterium extorquens* AM1) which was first isolated on methylamine (AM1 means Airborne Methylamine 1). 1968 -

He showed that oxidation is catalysed by a dye-linked methylamine dehydrogenase, assayed in a reaction system like methanol dehydrogenase.



The prosthetic group of methylamine dehydrogenase is TTQ, tryptophan tryptophyl quinone [McIntire et al 1991]. It is covalently bound, being a modified part of the protein backbone.

Remarkable fact: the first 2 types of quinoprotein to be described were both isolated in *Methylobacterium extorquens* AM1.


The electron acceptor for methylamine dehydrogenase was shown in 1981 by Tobari and Harada to be a blue copper protein called Amicyanin. This is a *single electron* acceptor.

The prosthetic groups of some Quinoproteins


The prosthetic group of methanol dehydrogenase is PQQ, Pyrrolo-quinoline quinone [Kennard's group 1979]

The prosthetic group of methylamine dehydrogenase is TTQ, tryptophan tryptophyl quinone [McIntire et al 1991]


The prosthetic group of the copper containing amine oxidases is TPQ, topa-quinone [Klinman's group 1990]



PQQ




TPQ




TTQ

Prosthetic groups of quinoproteins

Dehydrogenases




Not covalently bound




TTQ

[Derived from tryptophan]



CTQ

Oxidases




TPQ

LTQ

[derived from tyrosine]

PQQ, pyrroloquinoline quinone
 TTQ, Tryptophan tryptophylquinone
 CTQ, cysteine tryptophylquinone
 TPQ, Topaquinone
 LTQ, Lysine tyrosylquinone

Electron transport during methylamine oxidation




1985 - 1990 Ashley Lawton and Kevin Auton

[Amicyanin – *Tobari*]

Bob Eady & Peter Large [MeNDH; 1968-1971]

Electron transport during methylamine oxidation




1985 - 1990 Ashley Lawton and Kevin Auton

[Amicyanin – *Tobari*]

Bob Eady & Peter Large [MeNDH; 1968-1971]

Methylamine dehydrogenase [Bob Eady and Peter Large, 1968-1971]



Electron transport chain [Ashley Lawton & Kevin Auton, 1983-1989]



Ashley Lawton
President, Phyllos Inc.



Kevin Auton
Founder and CEO
of NextGen

Much of the subsequent study of the mechanism of methylamine dehydrogenase has been done by Victor Davidson and the genetics in Mary Lidstrom's group (with Mila Chistaserdova)



Other amine oxidising enzymes

Methylamine oxidase (Amine oxidase) In *Arthrobacter* P1




The peroxide is removed by catalase. No useable energy from this reaction.
Its prosthetic group is probably TPQ as in other copper-containing amine oxidases.

Systems involving methylated amino acids

In 'non-pigmented pseudomonads': *Ps. aminovorans*; *Pseudomonas* MA, & *Hyphomicrobium*

Two step system




The dehydrogenase is usually a flavoprotein that interacts like other flavoproteins with the electron transport chain at the level of ubiquinone and cytochrome b.
(There are some reports of NAD-linked dehydrogenases in some *Pseudomonas* and *Hyphomicrobia* Netrusov)

The next slide shows some alternative two-part systems

Some alternative two-part systems

Alternative systems for production of N-methylglutamate are in *Pseudomonas* MS and *Hyphomicrobium* but they may not be involved in growth. Kung & Wagner, Loginova & Trotsenko, Meiberg & Harder



Oxidation of tetramethylammonium salts to trimethylamine and formaldehyde

In Organism 5H2. Hampton & Zatman 1973. A non-pigmented Gram-negative, non-motile facultative methylotroph. No energy is available from this reaction

Mono-oxygenase



Oxidation of trimethylamine to dimethylamine and formaldehyde

In obligate methylotrophs; organism W3A1, & 4B6 (like *Methylophilus*), and *Hyphomicrobium* Colby & Zatman 1973, 1974

Trimethylamine dehydrogenase



The enzyme is an unusual flavoprotein, interacting with the electron transport chain by way of a second flavoprotein, cytochrome b etc. So provides energy

Indirect route for trimethylamine oxidation


A trimethylamine mono-oxygenase produces the N-oxide which is then demethylated to formaldehyde plus dimethylamine. During growth on trimethylamine. In *Pseudomonas aminovorans* (a typical *Pseudomonas* sp.) and *Bacillus* PM6. Boulton & Large; Myers and Zatman.

a) Trimethylamine mono-oxygenase. No energy available



b) Trimethylamine N-oxide demethylase No energy available

It is the 2nd enzyme for oxidation of trimethylamine and also the 1st enzyme for growth of *Bacillus* PM6 on trimethylamine N-oxide



The oxidation of dimethylamine to methylamine and formaldehyde

There are two types of enzyme; the same one operates when dimethylamine is a growth substrate as when it is an intermediate in the oxidation of higher methylated amines.

Dimethylamine mono-oxygenase No energy is available from this reaction

In *Pseudomonas aminovorans* (a typical pseudomonad). Bob Eady and Peter Large.

Also in *Hyphomicrobium* in aerobic conditions




Dimethylamine dehydrogenase Energy is available. Meiberg and Harder 1978.

In *Hyphomicrobium* during anaerobic growth with nitrate as electron acceptor.



Similar to trimethylamine dehydrogenase. An unusual flavoprotein that interacts with electron transport chain before cytochrome b.

Oxidation of methylated amines



(a) Tetramethylammonium monooxygenase; (b) trimethylamine dehydrogenase (in obligate methylotrophs); (c) trimethylamine monooxygenase (in facultative methylotrophs); (d) trimethylamine TV-oxide demethylase (aldolase); (e) dimethylamine monooxygenase (all methylotrophs); (f) dimethylamine dehydrogenase (anaerobic *Hyphomicrobia*); (g) amine oxidase (in *Arthrobacter* and methazotrophic yeasts); (h) methylamine dehydrogenase; (i) N-methylglutamate synthase; (j) and (k) N-methylglutamate dehydrogenases.

Distribution of routes for oxidation of methylated amines

Trimethylamine

NAD-independent TMN dehydrogenase: Obligate methylotrophs (eg *Methylophilus*) and *Hyphomicrobium*

Mono-oxygenase system: All other methylotrophs including *Bacillus*

Dimethylamine

Mono-oxygenase: All aerobic methylotrophs including *Hyphomicrobium* growing aerobically

Dimethylamine dehydrogenase: *Hyphomicrobium* growing anaerobically

Methylamine

There are three types of system.

Methylamine dehydrogenase (MD); Methylamine oxidase (MO); N-methylglutamate-linked (NMG)

Most obligate methylotrophs	MD
Facultative autotrophs	MD
Pink facultative methylotrophs	MD or NAD-dependent or NAD-independent NMG dehydrogenase
Non-pigmented pseudomonads	NAD-independent NMG dehydrogenase
<i>Hyphomicrobia</i>	NAD-dependent or NAD-independent NMG dehydrogenase
<i>Arthrobacter</i> and yeasts	MO

NOTE: Some bacteria may have more than one system but this has rarely been tested.

There is no correlation between the system for methylamine oxidation and assimilation pathway

[Top of oxidation of methylated amines](#)

[Top of summaries](#)

[The unusual cytochromes and electron transport in methylotrophs](#) [PPT](#) [Return to History](#) [Top of summaries](#)

The strange structures of special cytochromes c from methylophilic bacteria

The molecular structure of an unusual cytochrome c_2 determined at 2.0 Å; the cytochrome c_H from *Methylobacterium extorquens*

Jon Read, Raj Gill, Simon L. Dales, Jon B. Cooper, Steve P. Wood, and Chris Anthony *Protein Sci.* 1999 8: 1232-1240

The 1.6 Å X-ray structure of the unusual c-type cytochrome, cytochrome c_L , from the methylophilic bacterium *Methylobacterium extorquens*

Paul Williams, Leighton Coates, Fiyaz Mohammed, Raj Gill Peter Erskine, Dominique Bourgeois, Steve P. Wood, Chris Anthony and Jonathan B. Cooper*
J. Mol. Biol. (2006) 357, 151–162

INTRODUCTION:

cytochrome c: methylophilic: methanol oxidation: electron transport chains
methylophilic cytochromes c (earlier work)

Cytochrome c (Class I)

Mitochondrial cytochrome c and the equivalent bacterial cytochrome c₂.

Small water soluble protein; about 100 amino acids

Hundreds of sequences and many structures are known.


Haem prosthetic group covalently bonded by thioether bridges to protein


The iron atom forms ligands with the 4 pyrrole N of the haem; the 5th and 6th ligands are histidine and METHIONINE

Electron carriers $\text{Fe}^{+++} + \text{e}^- \longleftrightarrow \text{Fe}^{++}$


They function in electron transport chains, mediating electron transfer between membrane cytochrome bc₁ complex and the oxidase [or between membrane proteins in photosynthetic electron transport in bacteria]

They are located between the inner and outer membranes of mitochondria and in the equivalent periplasmic space in bacteria






Cytochrome c

Bacterial cytochrome c₂

Propionates




Haem

Iron

The planar haem lies enclosed in the haem cleft; electrons enter and leave the haem at its exposed edge

In all structures there is a water molecule in the LHS of the haem cleft; the movement of this molecule is said to be essential during the oxidation and reduction processes

Electron transport chain [Garrett & Grisham]



Electron transport chain in mitochondria






FIGURE 15.3

Respiratory electron carriers in the mitochondrion. This figure shows the sequence of electron carriers that oxidize succinate and NAD⁺-linked substrates in the inner membrane.



Text books are nearly all wrong

[Mathews]
Cyt c is a membrane protein!



Methylobacterium extorquens

- *Pseudomonas* AM1 (Peel & Quayle, 1961)
- *Pseudomonas* sp. M27 (Anthony & Zatman, 1964)



The oxidation of methanol

Catalysed by soluble Methanol dehydrogenase [MDH] in the periplasm. NAD is not involved.

MDH has a 'novel' prosthetic group called PQQ

MDH is equivalent to a soluble flavoprotein but with PQQ instead of FMN or FAD

The question: how is energy obtained from this oxidation of methanol; how does MDH interact with the electron transport chain?

Cytochrome c is particularly important in methanol oxidation [kinetic and mutant studies].

Biochem. J. (1980) **192**, 411–419
 Printed in Great Britain

The two cytochromes *c* in the facultative methylotroph *Pseudomonas* AM1

David T. O'KEEFFE and Christopher ANTHONY
 Department of Biochemistry, University of Southampton, Southampton SO9 3TU, U.K.

Cytochrome c_H High isoelectric point [basic, +ve charge at pH 7; $M_r = c\ 10,000$
 Does not react with MDH; **rapidly oxidised by the oxidase.**
 It is also involved in oxidation of NADH
 This cytochrome appears to be a typical bacterial cytochrome c_2


Cytochrome c_L Low isoelectric point [acidic, –ve charge at pH 7; $M_r = c\ 20,000$
Reacts rapidly with MDH; not oxidised by the oxidase.
 Not involved in NADH oxidation

Reacts slowly with CO. This displaces the 6th Methionine ligand.
 This is very rare in cytochromes *c* and indicates an exceptionally flexible structure in the LHS of the haem cleft

The two cytochromes *c* are able to react very rapidly with each other

MeOH $\rightarrow$ MDH $\rightarrow$ Cyt c_L $\rightarrow$ Cyt c_H $\rightarrow$ oxidase $\rightarrow$ O₂

The oxidation of methanol in methylotrophs



Project: Solve structure of all the soluble proteins

Study electron transport within the proteins and between them

The path of electrons and protons during the oxidative part of the reaction mechanism of MDH


Electrons must pass from reduced PQQ (PQQH₂) to cytochrome c_L

Cytochrome c_L


Note involvement of the disulphide ring

Protons must pass from PQQH₂ by way of its hydroxyls, and the hydrogen-bonding network, to the water (W904) which is accessible to the periplasm

The pathways are based on those proposed for the quinoxaemoprotein alcohol dehydrogenase by Oubrie, Dijkstra *et al*, and Chen, Matsushita, Toyama, Scott Mathews *et al*.



Interaction of the electron transport proteins




MDH and cytochrome c_L 'dock' together by ionic interactions between lysines on MDH and carboxylates on cytochrome c_L.

The electron transfer complex may involve interaction of the cytochrome with the hydrophobic funnel on the surface of MDH.

Cytochrome c_L dissociates, and then passes electrons to cytochrome c_H.

Cytochrome c_H is almost a typical cytochrome c, passing its electrons to the oxidase



Claude Chan, Darren Day, Jon Cox, Jim Elliott, Simon Dales



Biochem. J. (1980) **192**, 411–419
Printed in Great Britain

The two cytochromes *c* in the facultative methylotroph *Pseudomonas* AM1

David T. O'KEEFFE and Christopher ANTHONY

Department of Biochemistry, University of Southampton, Southampton SO9 3TU, U.K.

Cytochrome c_H High isoelectric point [basic, +ve charge at pH 7; $M_r = c$ 10,000

Cytochrome c_L Low isoelectric point [acidic –ve charge at pH 7; $M_r = c$ 20,000




Fig. 5. pH-dependence of the midpoint redox potentials of cytochromes c_H and c_L .


Details of the methods are given in the Materials and methods section. Oxidative and reductive titrations gave identical results. ●, Midpoint potentials of cytochrome c_H ; ■, midpoint potentials of cytochrome c_L . The dotted lines are theoretical curves calculated from the equation:

$$E_m = E_0 + 59 \log \left(\frac{[H^+]^2 + K_{a1}[H^+] + K_{a1}K_{a2}}{[H^+]^2 + K_{r1}[H^+] + K_{r1}K_{r2}} \right)$$

where K_{a1} and K_{a2} are the dissociation constants for the first and second ionizations in the oxidized species and K_{r1} and K_{r2} are the equivalent dissociation constants for the reduced species. The pK ($-\log K$) and E_0 values assumed for calculation of the theoretical curves are given in Table 3.

There were four ionizations affecting the redox potentials in the pH range studied (pH 4.0–9.5), two in the oxidized form (pK values about 3.5 and 5.5) and two in the reduced form (pK values about 4.5 and 6.5),

suggesting that the ionizing groups involved may be the two propionate side chains of the haem.



This suggests the propionates are very unusual in being exposed to solvent and not buried within the cytochrome structure as with other cytochromes


Fig. 1. The sequence of cytochrome c_H of *M. extorquens*.

	-5	1	10	20	30	40	50
Yeast		TEFKAGSAKKGATLFKTRCLQCHTVEKGGPHKVGPNLHGI	FGRHSGQAE-GYST	YTDA			
Horse		GDVEKGKKIFVQKCAQCHTVEKGGKHKTGPNLHGLFGRKTGQAP-GF-TY	TDA				
Tuna		GDVAKGKKTFFVQKCAQCHTVENGKKHKVGPNLWGLFGRKTGQAE-GY-SY	TDA				
	1	10	20	30	40		
Methylobacterium		EGDAAAGEKAFA-PCKACHNFEKNG---	VGPTLKGVVGA	KAGEGADGYA-FSDA			
		60	70	80	90	112	
Yeast		NIK---KNVLWDENNMS	YLTNP-KYIPG-----	TKMAFAGLKKEKDRNDITYLKKACE			
Horse		NKN---KGITWKEETLMEY	LENPKKYIPG-----	TKMIFAGIKKKTEREDLIAYLKATNE			
Tuna		NKS---KGIVWNENTLMEY	LENPKKYIPG-----	TKMIFAGIKKKGERQDLVAYLKSATS			
	50	60	70	80	90	100	
Methylobacterium		LKK---SGLTWDQADLK	WLADEPKKKVPG-----	TKMVFPGISDPKKVDDIIAYLKTKS			


About 50% identity with sequence of typical cytochromes c

Almost all 'conserved residues' are present

Residues in red box are always Tyrosine
(100% conservation in 160 sequences).
But replaced by tryptophan in cytochrome c_H



Tyrosine



Tryptophan


The molecular structure of an unusual cytochrome c_2
determined at 2.0 Å; the cytochrome c_H from
Methylobacterium extorquens

JON READ, RAJ GILL, SIMON L. DALES, JON B. COOPER, STEVE P. WOOD,
AND CHRIS ANTHONY

Division of Biochemistry and Molecular Biology, School of Biological Sciences, University of Southampton,
Southampton SO16 7PX, United Kingdom


Protein Science (1999), 8:1232–1240.

The overall folds of cytochrome c_H , cytochrome c , and cytochrome c_2 .




Cytochrome c_H is unique in being a bacterial cytochrome c that is folded into a mitochondrial cytochrome c structure

The exposed propionates in cytochrome c_H




The LHS of the haem cleft in the cytochromes c

cytochrome *c*



cytochrome *c_H*



The 'absolutely conserved' Tyr67 is replaced by tryptophan

The 'conserved' water is absent; the LHS of haem cleft is hydrophobic

This shows that the previously proposed mechanism of cytochrome c, involving the 'essential water' is no longer valid.

The 1.6 Å X-ray Structure of the Unusual c-type Cytochrome, Cytochrome c_L , from the Methylophilic Bacterium *Methylobacterium extorquens*

Paul Williams¹, Leighton Coates¹, Fiyaz Mohammed¹, Raj Gill¹
Peter Erskine¹, Dominique Bourgeois², Steve P. Wood¹
Chris Anthony¹ and Jonathan B. Cooper^{1*}

¹School of Biological Sciences
University of Southampton
Bassett Crescent East
Southampton SO16 7PX

J. Mol. Biol. (2006) **357**, 151–162

172 residues instead of usual 100. Has 40-50 extra residues at N terminus
and a C-terminal extension

Negligible sequence identity to other cytochromes c.


Predictions:

The haem propionates would be exposed to solvent

The loop bearing the 6th ligand methionine might be unusually flexible

The two very unusual cysteines would be involved in a disulphide bridge to hold the long loops together in the structure

The structure of cytochrome c_L



172 amino acids

No sequence similarity


Similar basic structure
with additional features

Paul Williams

Cytochrome c_L structure

Helices A, C and E are
usual haem-enclosing
helices

His112 has replaced Met
109 as ligand to haem iron;
Loop 3 is relatively flexible
in this cytochrome leading
to reaction with CO







Figure 4. Surface view of cytochrome cL. The outer haem propionate (HP6) is clearly exposed to solvent as are the two waters (W6 and W7). W7 is bonded to the inner haem propionate (HP7). Asp83, one of the ligands to the Ca^{2+} ion, has been proposed to be involved in interaction with methanol dehydrogenase together with Glu120, the outer haem propionate (HP6) and those residues coloured green²⁰.

Figure 4. The calcium ion and the interactions of the two haem propionates, in cytochrome c_L .



HP6 is the outer haem propionate and HP7 is the inner haem propionate. The lines indicate hydrogen bonds or coordinations. Water molecules (Wat6–Wat9, Wat115, and Wat127) are indicated by blue spheres; the calcium ion is indicated by a magenta sphere; the haem iron is indicated by a red sphere. The calcium ion and the oxygen atoms of the four water molecules (Wat6–Wat9) and the side-chain of Asp83 lie in one plane; the bonds to the oxygen atoms of Gly80 and Tyr85 (shown as continuous lines) are perpendicular to this plane. In addition to the bonds indicated here, the main-chain carbonyl group of Asp83 is also hydrogen bonded to the side-chain nitrogen atom of Trp86, (not shown). The water molecules Wat6, Wat7 and Wat115 are exposed to bulk solvent (Figure 7).

Two unusual c-type cytochromes are involved in methanol oxidation pathway

Cytochrome c_H

Electron donor to the oxidase; Same sequence and function as typical cytochrome c

Typical cytochrome c structure except the folds are like mitochondrial (not bacterial) cytochrome

The 'conserved' tyrosine and water are absent; the hydrophobic haem cleft shows the water is not important in cytochrome mechanism

The propionates are exposed to solvent

Cytochrome c_L

Completely novel cytochrome. Although the 3-helix core is same as other cytochromes, there is negligible sequence identity and it is unique in lacking one of the 7 conserved positions found in all families of cytochromes c [haem binding and residues that hold 3 helices together in position]

A calcium ion replaces the 'conserved' arginine in balancing propionate charge

As predicted the haem propionates are exposed and there is a novel disulphide bridge

The two cytochromes, in the same organism and the same electron transport chain seem to have evolved completely independently

Acknowledgements:

David O’Keeffe, Mathew Beardmore-Gray, Sandy Cross, Jim Elliott, Bill Keevil, Sasha Netrusov, Claude Chan, Darren Day, Jon Cox, Simon Dales, Karen Amaratunga, Paul Skipp, Tosin Majekodunmi, Remi Afolabi,

Jon Read, Raj Gill, Jon B. Cooper, Steve P. Wood, Paul Williams, Leighton Coates, Fiyaz Mohammed, Raj Gill, Peter Erskine, Dominique Bourgeois,

Prof Akhtar for encouragement (“We are not interested in cytochromes, are we?”)

[Top of cytochromes](#)

[Top of summaries](#)

[The Methylo](#)

[PPT](#)

[Top of summaries](#)

[Return to History](#)

The Methylootrophs

They use compounds with one or more carbon atoms but no C-C bonds as their sole source of carbon and energy: Methane, Methanol, methylamine, dimethylamine.

Those that use methane are also called **Methanotrophs** (eg. *Methylococcus*, *Methylosinus*)

Obligate methylootrophs cannot grow on multicarbon compounds (include most methanotrophs)

Facultative methylootrophs also grow on multicarbon compounds (eg. *Methylobacterium extorquens* = *Pseudomonas* AM1)

Restricted facultative methylootrophs grow on C1 or C2 compounds (eg. *Hyphomicrobium*)

Facultative autotrophs grow on carbon dioxide, plus hydrogen or methanol by way of RuBP path (eg *Paracoccus*)

There is a huge number of different types of methylootroph. Very few are typical bacteria that had been described previously.

Although their oxidative metabolism is basically similar, there are many different carbon assimilation pathways.


Most are Gram-negative. The few Gram positive methylootrophs (*Bacillus* sp) have different oxidation enzymes

- 1906 Sohngen. *Bacillus methanicus* growing on methane
- 1961 Peel and Quayle. *Pseudomonas* AM1 (cited only 6 papers) A pink facultative methylootroph
- 1964 Anthony and Zatman. Description of Methanol dehydrogenase in *Pseudomonas* M27 (=AM1)
- 1964 Hirsch and Conti. *Hyphomicrobium*. A stalked facultative methylootroph
- 1970 Whittenbury (with Wilkinson). 100 new methanotrophs (2 Types)
- 1971 Ribbons Early work on methane oxidation
- 1970- Ogata, Sahm, Hazeu, van Dijken, Harder. Methylootrophic yeast.
- 1980 Dalton Definitive work on methane oxidation
- 1982 Anthony book. Cited about 1200 methylootroph references

The main 'workhorses' in the study of methylotrophs

- *Pseudomonas* AM1 (Peel & Quayle, 1961)
- *Pseudomonas* sp. M27 (Anthony & Zatman, 1964)

Methylobacterium extorquens. A pink facultative methylotroph



Me in 1963



J Rod Quayle



My PhD supervisor and my first research student (Len Zatman and Pat Dunstan, now Goodwin)




Howard Dalton, Rod and me


Methylococcus capsulatus (Type I obligate methanotroph)
Methylosinus trichosporium (Type II obligate methanotroph)



Type I membranes




Type II membranes



Doug Ribbons 1973




Rod Quayle & Roger Whittenbury



John Wilkinson

Hyphomicrobium



Sticks to surfaces; grows only on C1 and C2 compounds. Reproduce by budding

Grow on methanol and methylated amines but not methane

Hirsch & Conti, Harder, Dow, Attwood.
Quayle (assimilation pathways)



Hans van Dijken and Wim Harder 1973
Hyphomicrobium, Yeast and Growth
yields of methylotrophs

Methylotrophic Yeasts


By contrast with bacteria, the methylotrophic yeasts are not a special group; they are in the usual genera of yeasts [*Hansenula*, *Candida*, *Pichia* and *Torulopsis*]. Only 7/39 genera grow methylotrophically. They grow only on methanol (not methylamines or methane). Again, by contrast with bacteria they all share the same assimilation pathways.

They oxidise methanol by a flavoprotein alcohol oxidase coupled to catalase which is present in crystalline form in peroxisomes and which yields no energy. Almost all bacteria use an energy-yielding quinoprotein methanol dehydrogenase. all


1969-1975

Ogata, Tani, Sahm, van
Dijken, Harder,
Quayle (assimilation)

Peroxisomes in methanol-oxidising yeast




The oxidase is produced in very large amounts and transcription of the gene has a very strong promoter; it is now used in genetic engineering for production of large amounts of cloned Eucaryotic proteins.



Crystals of the flavoprotein alcohol oxidase in yeast

Methanol oxidation in methylotrophic yeast

This is catalysed by non-specific flavoprotein called alcohol oxidase
This is in a peroxisome with catalase



The enzyme is used as a source of hydrogen peroxide –
for wound disinfection and also used in washing powder



J. Guis Kuenen and Professor Kato (Tottori and Kyoto)



Friends from Yamaguchi, Japan: Matsushita, Uragami (?), Ameyama, Adachi

The main 'groups' of bacterial methylotrophs

Methanotrophs. Vast majority are obligate. (Recent important exceptions)

All have internal membranes and produce spores. Gram negative.

Type I (eg *Methylococcus*, *Methylomonas*) use RuMP pathway

Type II (eg *Methylosinus*, *Methylocystis*) use Serine pathway

Methylotrophs not using methane. None have internal membranes or produce spores. Vast majority are Gram negative.

Obligate methylotrophs Grow on methanol and/or methylamine. Gram negative, motile, non-sporing. Use RuMP pathway. Eg *Methylophilus*

Facultative methylotrophs. Usually grow on methanol/methylamine; use serine pathway

Pink facultative methylotrophs. Motile, single polar flagellum. Red carotenoids similar to photosynthetic bacteria. Wide range of C substrates but few carbohydrates. *Methylobacterium*

Non-pigmented facultative 'pseudomonads'. No typical pseudomonas sp use C1 compounds.

Pseudomonas aminovorans. Use serine pathway (icl⁺ variant)


Gram-negative or variable non-motile rods. Some only use methylated amines. A mixed group. Eg *Arthrobacter*.

Facultative autotrophs or phototrophs Use RuBP pathway (Calvin cycle). *Paracoccus denitrificans* (can use hydrogen); *Rhodopseudomonas*

Gram positive facultative methylotrophs Very rare. *Bacillus*, *Streptomyces*

Hyphomicrobia Restricted facultative methylotrophs. Stalked. Budding. Serine pathway

The ICI 'Pruteen' plant at Billingham, UK 1980



The centre tower is the 1.5 million litre fermenter
It contains about 100,000,000,000,000,000,000 methylotrophs
10% of their soluble protein is Methanol dehydrogenase

The Methylootrophs and their biochemistry as was known up to 1981 is covered comprehensively in


The Biochemistry of Methylootrophs by Chris Anthony

It can be downloaded complete and free as a pdf file from my website

www.chris-anthony.co.uk

On this page: <http://www.chris-anthony.co.uk/methylootrophs.html>

The Biochemistry of Methylootrophs 1982



The Biochemistry of Methylootrophs

C. ANTHONY

*Department of Biochemistry
University of Southampton
England*

Chapters 12

Text pages 350

Index 48 pages

Tables 60

Figures 57

References 1300

1982



ACADEMIC PRESS

A Subsidiary of Harcourt Brace Jovanovich, Publishers

London · New York
Paris · San Diego · San Francisco
São Paulo · Sydney · Tokyo · Toronto

Methylootrophic bacteria
Ribulose biphosphate pathway
Ribulose monophosphate cycle
The Serine pathway
The TCA cycle and growth
on multicarbon compounds
Oxidation of methane, methanol,
formaldehyde and formate
Oxidation of methylated amines
Electron transport and
energy transduction
Growth yields and bioenergetics
Methylootrophic yeasts
Methanogens and
methanogenesis
Commercial exploitation

A CD version is available and the complete book
is on my website: chris-anthony.co.uk


The Biochemistry of Methylotrophs

C. Anthony (1982) Academic Press, London

This textbook is no longer in print.

It contains 350 pages, 60 pages of Tables, 60 pages of Figures, a 40 page index and about 1000 references [every methylotroph reference up to 1981]. Copyright exclusive to author. Contents are summarised [below](#).

I have produced a CD of the book with all pages in both jpg and pdf formats which can be searched in Acrobat Reader. A copy of the CD is available [£25 or \$50]. Email: C.Anthony@soton.ac.uk




For complete book click on picture

Dedicated to K. G. Wiles, L. J. Zatman and J. R. Quayle, FRS

Foreword by Professor J.R. Quayle FRS

There are two kinds of organism which stand out as the biosynthetic virtuosi of the living world—the autotrophs and the methylotrophs. The autotrophs are organisms which can synthesize all their cell constituents from carbon dioxide as sole carbon source; methylotrophs can perform such total biosynthesis from reduced carbon compounds containing one or more carbon atoms but containing no carbon-carbon bonds. Autotrophs have long been the subject of



Click on book ot download it complete (free). Or ask for CD (free)

[Top of Methylotrophs](#)
[Top of Summaries](#)

[Growth yields of Methylotrophs](#)

[PPT](#)

[Top of Summaries](#)

Factors affecting growth yields in methylotrophs

For growth on a simple defined medium with a single source of carbon:

Growth yield is Grams dry weight cells / moles of growth substrate consumed

$Y_s = \text{g/mole}$

The substrate is consumed for 2 main purposes: production of new cells plus provision of energy (ATP and NADH) for the necessary biosynthesis.

In anaerobes: At least 95% of the substrate is used for energy (mainly ATP) production. This is produced by fermentation where ATP is produced by substrate level phosphorylation.

The yield is determined by the number of ATPs produced per mole of substrate.

In conventional aerobes: About 50% of the carbon substrate is used for energy production. This substrate is oxidised completely to CO_2 and the ATP is produced by oxidative phosphorylation. *****

Yields are expressed as $Y_s = \text{g/mole substrate used}$ and $Y_{\text{O}_2} = \text{g/mole of O}_2$

Yields are dependent on the P/O ratio = moles ATP produced / atom of oxygen

consumed during oxidative phosphorylation. = moles ATP / mole of NADH oxidised


Yield per mole of ATP consumed is $Y_{\text{ATP}} = \text{g/mole ATP}$

This is often (justifiably) considered to be a constant.


This assumes that the ATP requirement for transport and biosynthesis is similar for all substrates.

Oxidation of organic carbon substrates

a) Anaerobic oxidation (Fermentation)



b) Aerobic oxidation



Note: methylotrophs are special as the substrates are oxidised directly to CO₂

Yield per mole of ATP consumed is $Y_{ATP} = g/\text{mole ATP}$

This is often (justifiably) considered to be a constant. It assumes that the ATP requirement for transport and biosynthesis is similar for all substrates and that the growth yield is directly proportional to the ATP available. Such microbes (the vast majority) are said to be ATP-limited

Growth yield predictions and measurements in methylotrophs illuminate those aspects of their biochemistry and physiology that make them special. It is important when considering C1 compounds as a substrate for methylotrophs as a source of Single Cell Protein, SCP. Also when considering biomass production in ecological studies.

Most multicarbon substrates are at the level of oxidation of CH_2O [glucose, formaldehyde etc]

It might be assumed that more reduced substrates (hydrocarbons, long chain fatty acids, methane, methanol, methylamine), having more energy would give high yields. **Not true.**

To be true then all of this energy must be harnessed as NADH and ATP during oxidation.

The Truth: Alkanes are chemically inert and use energy (as NADH) in the initial hydroxylation step. So they are equivalent to CH_2O .

The oxidation of methanol and methylamine use unusual enzymes that are not NAD-linked and yield relatively little ATP in their electron transport chains.

The prediction of growth yields

- a) Substrate is converted to a central precursor (eg phosphoglycerate, PGA). This also produces some NADH and ATP
- b) PGA is biosynthesised into cell material. This requires NADH and ATP
- c) More substrate is oxidised to produce the NADH for this biosynthesis
- d) More substrate is oxidised to produce ATP for biosynthesis

This can be expressed in equations which lead to an overall Assimilation equation. This can be used to investigate the effects of different assimilation pathways and energy production systems.

- a) Calculations for a typical multicarbon substrate, glucose
- b) The results of similar equations for methylotrophs

Much of the following is taken direct from a paper on prediction of growth yields and a chapter in *The Biochemistry of Methylotrophs*

Assumptions: the constituents of most cells are similar as are the pathways for their biosynthesis. It is assumed that the nitrogen source is ammonia.

The Assimilation equation for glucose

The assumptions listed above lead to the following equation for assimilation of phosphoglycerate (PGA) into cell material (in these equations H^+ , H_2O , NAD^+ , ADP and P_i are omitted for clarity):



By taking into consideration the biosynthesis of PGA from growth substrate, an overall assimilation equation may be derived.


For example, if the formula for glucose transport and metabolism to PGA is:




then the assimilation equation for glucose will be:




The NADH is produced by oxidising glucose to CO_2 (FPH₂ is reduced succinate dehydrogenase):



By adding the last two equations, the overall assimilation equation (simplified) for predicting the effect of varying P/O ratios on growth yields on glucose is obtained:



Glucose assimilation equation

If glucose is metabolised by glycolysis and TCA cycle, and oxidation of NADH yields 3ATP then 1 glucose yields 38 ATP and 6 CO₂. So 0.8 glucose must be oxidised to give the ATP and also 4.8 CO₂

SO yield equation: 2.95 glucose gives 306 g cell + 5.7 CO₂ Yield = 104g/mole glucose

If the P/O ratio is only 2 instead of 3 only get 2ATP per mole NADH

For provision of 30.6 ATP need to oxidise 1.2 glucose, producing 7.2 CO₂

SO yield equation: 3.35 glucose gives 306 cell + 8.1 CO₂ Yield = 91 g /mole glucose

If P/O ratio is only 1 then 2.4 moles glucose is needed for ATP

Yield equation 4.55 glucose gives 306g cell + 15.3 CO₂ Yield = 67g / mole glucose

Carbon conversion efficiency if P/O ratio is 3 is 306 g/540g glucose = 57%

If P/O ratio is only 2 then CCE is 51%

If P/O ratio is only 1 then CCE is 37%

Bacteria growing on glucose are ATP- limited

Prediction of growth yields in methylotrophs

a) Substrate is converted to a central precursor (eg phosphoglycerate, PGA). This usually produces some NADH and ATP with multicarbon substrates

In methylotrophs this first part involves initial oxidation to formaldehyde. With methane this **uses** oxygen and NADH to oxidise the methane to methanol. Oxidation of Methanol (and methylamine) to formaldehyde does not produce NADH but only reduced quinoprotein dehydrogenases that yield only 1ATP (or less) during their oxidation.

Note: The special pathways for conversion of formaldehyde to PGA sometimes consume a lot of NADH (RuBP and Serine pathways)

b) PGA is biosynthesised into cell material. This requires NADH and ATP as in all bacteria

c) More substrate is oxidised to produce the NADH for this biosynthesis.

Again there is a problem: The first step in methane oxidation **uses** NADH. The oxidation of methanol and methylamine to formaldehyde produces no NADH. This is only produced during formaldehyde oxidation (not always) and formate oxidation (essential).




d) More substrate is oxidised to produce ATP for biosynthesis

During oxidation of methanol or methylamine to provide NADH the reduced enzymes were re-oxidised by the electron transport chains, giving ATP. Because of this it is often the case that no substrate needs to be oxidised for this final provision of ATP

NADH limitation and growth yields

Anthony, 1978



A: MeOH is oxidised to HCHO and assimilated into CELL using ATP and NADH.

B: MeOH is oxidised to provide NADH for assimilation pathway.

C: The reduced MDH produced in A and B is oxidised to give ATP.

D: MeOH is oxidised to produce any ATP that is still required.

During oxidation of methanol so much ATP is produced in step C that little further ATP production is required. The cells are NADH-limited instead of ATP-limited. The growth yield is not dependent on the ATP yield from methanol dehydrogenase.


On methane: the first step requires NADH and so the effect is even more marked.

Assimilation equations for the three main methylotroph pathways



Ribulose biphosphate pathway:



Ribulose monophosphate pathway:



Serine pathway:



RuBP and Serine pathways need a lot of NADH; RuMP pathway needs relatively little

Using the equations the effects of the following characteristics on growth yields can be evaluated:

P/O ratio

Hydroxylation of methane and methylated amines using NADH

The assimilation pathway

The nature of the system for oxidation of formaldehyde (yielding NADH or not)


Conclusions:

Most methanotrophs are NADH-limited regardless of assimilation pathway

On methanol, bacteria with the RuMP pathway are sometimes typically ATP-limited but changing the P/O ratio will have little effect on growth yield (they tend to be carbon-limited, where the only way of increasing yield is to add oxidised substrates).

Most serine pathway bacteria will be NADH-limited where a different P/O ratio will have little effect on yield

The ICI 'Pruteen' plant at Billingham, UK 1980



The centre tower is the 1.5 million litre fermenter

It contains about 100,000,000,000,000,000,000 methylotrophs
10% of their soluble protein is Methanol dehydrogenase

[Top of Growth yields](#)
[Top of Summaries](#)

[Carbon assimilation pathways](#)

[PPT](#)

[Top of Summaries](#)

[Return to History](#)

Carbon assimilation pathways

Part one: Brief summary of the four pathways for assimilation of C1 compounds

The elucidation of the Serine Cycle up to 1973

Part two: The solution of the complete Serine / Ethylmalonyl-CoA cycle

Gordon Research Conference: Magdalen College, Oxford, 2006

Molecular Basis of Microbial One-Carbon Metabolism

The Biochemistry of Methylotrophs: a historical perspective

Chris Anthony, University of Southampton, UK



Dedicated to the memory of J. Rod Quayle
(1926 – 2006)

- 1946 – 1951 PhD in physical-organic chemistry
[University of Wales; with ED Hughes]
PhD on aphid pigments [Cambridge with
Alexander Todd]
- 1953 – 1954 Calvin's lab at Berkeley
- 1955 – 1963 Krebs' MRC Unit at Oxford
- 1963 – 1983 University of Sheffield
- 1983 – 1992 Vice-Chancellor, University of Bath

Many of my slides are from this lecture
dedicated to Rod Quayle

Carbon assimilation pathways of methylotrophs

Ribulose biphosphate [RuBP] pathway [Key contribution from JRQ]

Plants, autotrophic bacteria and a few methylotrophs

Pathways first proposed by Quayle and mainly elucidated by him and his colleagues:

Ribulose monophosphate [RuMP] pathway

Type I methanotrophs and obligate methanol or methylamine utilisers

Dihydroxyacetone [DHA] pathway

Methylotrophic yeasts


Serine pathway

Type II methanotrophs and facultative methanol or methylamine utilisers

I will summarise the first three and spend more time on details of serine pathway

Calvin-Benson cycle for CO₂ fixation in plants [1950 – 1960]

The key demonstration of the specific RuBP carboxylase activity in extracts was published by Quayle in JACS in 1954



RADIATION LABORATORY AND
DEPARTMENT OF CHEMISTRY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA


RECEIVED MAY 24, 1954

J. R. QUAYLE
R. C. FULLER
A. A. BENSON
M. CALVIN

This pathway was soon shown to be the path of carbon dioxide fixation in aerobic autotrophic bacteria and it was commonly assumed that methylotrophs growing on methane or methanol would assimilate their carbon by this pathway after their oxidation to CO₂

JRQ showed that this is the route for formate assimilation by *Pseudomonas oxalaticus* [1959].

He later showed that the facultative autotroph *Paracoccus denitrificans* assimilates methanol by this pathway.



Ribulose Bisophosphate pathway in plants, autotrophs and some methylotrophs



Figure 1. Some members of the photosynthesis research group (about 1953). They are in front of Old Radiation Laboratory at UCB. This building was later demolished to make space for a new Department of Chemistry building, Hildebrand Hall. *Left to right:* Clinton Fuller, Robert Norris, Hans Kornberg, Alice Holtham, Melvin Calvin, J. Rodney Quayle, Malcolm Thain, James Bassham and Jaques Mayudon. Andrew Benson agrees that he probably took this picture, accounting for his absence from the picture.

The ribulose monophosphate pathways

Quayle, Johnson, Strom, Ferenci, Kemp, [1965 – 1974]

Occur in Type I methanotrophs and in the obligate methanol or methylamine utilisers. There are 4 variants; three of these have been demonstrated in different bacteria.

Methods: Short term labelling experiments; analysis of position of label in metabolites, purification and characterisation of enzymes; measurement of all enzymes of the pathway.

Similar to Ribulose biphosphate (Calvin) cycle except for 'first reaction'

Condensation of formaldehyde with RuBP to give a novel hexulose phosphate; this is then isomerised to fructose 6 phosphate. The novel synthase and isomerase were isolated and characterised.

Subsequent reactions of the pathway are similar to the rearrangement reactions of the Calvin cycle.





Fig. 6. The ribulose monophosphate (RuMP) cycle of formaldehyde assimilation (KDPG aldolase/transaldolase variant). This variant occurs predominantly in obligate methylotrophs. The enzyme numbers refer to those listed in the text: (8) transketolase; (9) pentose phosphate epimerase; (11) pentose phosphate isomerase; (12) transaldolase; (13) hexulose phosphate synthase; (14) hexulose phosphate isomerase; (16) glucose phosphate isomerase; (17) glucose phosphate dehydrogenase; (18) phosphogluconate dehydrase; (19) 2-keto, 3-deoxy, 6-phosphogluconate (KDPG) aldolase; (20) PEP synthetase or equivalent enzyme(s); (21) enolase; (22) phosphoglyceromutase.






Fig. 8. The ribulose monophosphate (RuMP) cycle of formaldehyde assimilation (Fructose biphosphate aldolase/transaldolase variant). This variant does not occur frequently in methylotrophs. The enzyme numbers refer to those listed in the text: (2) phosphoglycerate kinase; (3) glyceraldehyde phosphate dehydrogenase; (4) triose phosphate isomerase; (5) aldolase; (8) transketolase; (9) pentose phosphate epimerase; (11) pentose phosphate isomerase; (12) transaldolase; (13) hexulose phosphate synthase; (14) hexulose phosphate isomerase; (15) phosphofructokinase.

The dihydroxyacetone [DHA] cycle of formaldehyde assimilation in yeasts

Nobuo Kato, O'Connor (Mary Lidstrom), Sahm, Babel, van Dijken, Quayle
[1977-1981]

This is similar to the RuBP and RuMP cycles

Two specific enzymes are required for formaldehyde fixation:

DHA synthase and triokinase


These were purified and characterised

Short term labelling pattern from ^{14}C methanol was consistent with the cycle proposed by Quayle and distribution of labelled carbon in the proposed intermediates was consistent with the cycle

Mutants lacking the key enzymes were unable to grow on methanol

DHA cycle in yeast

Fixation: xylulose phosphate + HCHO →
glyceraldehyde phosphate + dihydroxyacetone



The discovery of the Serine cycle in *Methylobacterium extorquens* AM1 (*Pseudomonas* AM1)

Part 1: J Rod Quayle, David Peel, Peter Large, Alan Salem
(with bits by Pat Goodwin (nee Dunstan) and Chris Anthony)





J. Rod Quayle



Peter Large

Methylobacterium extorquens

- *Pseudomonas* AM1 (Peel & Quayle, 1961)
- *Pseudomonas* sp. M27 (Anthony & Zatman, 1964)



The Serine Pathway; Peter Large, David Peel and Rod Quayle


1961 - 1963

1. Large, P.J., Peel, D. and Quayle, J.R. *Biochemical Journal* **81**, 470-480 (1961).
Microbial growth on C1 compounds: Synthesis of cell constituents by methanol- and formate-grown *Pseudomonas* AM1 and methanol-grown *Hyphomicrobium vulgare*.
2. Large, P.J., Peel, D. and Quayle, J.R. *Biochemical Journal* **82**, 483-488 (1962).
Microbial growth on C1 compounds: Distribution of radioactivity in metabolites of methanol-grown *Pseudomonas* AM1 after incubation with [14C]methanol and [14C]bicarbonate.
3. Large, P.J., Peel, D. and Quayle, J.R. *Biochemical Journal* **85**, 243-250 (1962).
Microbial growth on C1 compounds: Carboxylation of phosphoenolpyruvate in methanol-grown *Pseudomonas* AM1.
4. Large, P.J. and Quayle, J.R. *Biochemical Journal* **87**, 386-396 (1963).
Microbial growth on C1 compounds: Enzyme activities in extracts of *Pseudomonas* AM1.

The Elucidation of the Serine pathway in *Pseudomonas* AM1 [now *Methylobacterium extorquens* AM1]

A pink facultative methylotroph; grows on methanol, not methane

Bacteria were grown on ^{14}C MeOH and the label in cell material recorded. If RuBP pathway is operating then passage of 'cold' $^{14}\text{CO}_2$ would decrease the label by 95%



Passage of 'cold' CO_2 through the culture during growth on $^{14}\text{CH}_3\text{OH}$ decreased label in cell material by about 50%. This shows that half the carbon enters the biosynthetic pathway as CO_2 produced from the methanol

RuBP carboxylase is absent

Short term labelling experiments showed that 3- phosphoglycerate is not an early intermediate when whole cells are incubated with $^{14}\text{CH}_3\text{OH}$ or H^{14}COOH

Short term label experiments to determine path of carbon

Incubate growing cells with $^{14}\text{CH}_3\text{OH}$ or $^{14}\text{CO}_2$ (bicarbonate)

Take samples into boiling ethanol at 2,4,8,20 secs etc

Separate all soluble components by 2-way paper chromatography

Identify labelled compounds by autoradiography (3 weeks)

Elute, count ^{14}C and confirm identity by co-chromatography with known compounds

Plot % radioactivity in each compound against time. A negative slope indicates an early intermediate.

After 1 min incubation the early intermediates were chemically analysed to determine the specific radioactivity in each carbon atom

Distribution of label in cells incubated with labelled CO_2

Negative slope = earliest intermediates
 Malate [reflecting oxaloacetate, OAA]
 Glycine; Later - serine

Suggests typical carboxylation of a C3 to a C4 compound [OAA / malate]

And either cleavage of C4 to glycine

Or novel carboxylation to give glycine

NB: the presence of a labelled compound at 20 seconds does not indicate an early intermediate.

Coenzyme A derivatives are cannot be seen in this sort of experiment.




Fig. 3. Variation with time of the percentage distribution of ^{14}C incorporated from $[^{14}\text{C}]$ bicarbonate into the constituents of the ethanol-soluble fraction of *Pseudomonas* AM1 growing on methanol. ●, Malate; ▲, serine; □, glycine; ▼, phosphorylated compounds; x, trehalose.

Similar results were obtained using *Hyphomicrobium vulgare*

Distribution of label in cells incubated with methanol

Negative slope = early intermediates

Serine Malate Aspartate Glycine

Suggests:

Addition of HCHO to glycine to give serine

A derivative of serine is carboxylated to OAA / malate / aspartate




Fig. 2. Variation with time of the percentage distribution of ^{14}C incorporated from $[^{14}\text{C}]$ methanol into the constituents of the ethanol-soluble fraction of *Pseudomonas* AM1, growing on methanol. ●, Malate; △, aspartate; ▲, serine; □, glycine; ■, glutamate; ▼, phosphorylated compounds; ×, trehalose.

Similar results were obtained using *Hyphomicrobium vulgare*

Distribution of ^{14}C in carbon atoms of early intermediates


Cells were incubated for 1 minute with ^{14}C MeOH or $^{14}\text{HCO}_3$; Intermediates were purified, chemically degraded and ^{14}C in each C atom determined and expressed as % of total counts in the compound

	From methanol	From bicarbonate
Glycine		
CH_2NH_2	50	15
COOH	50	85
Serine		
CH_2OH	50	2
CHNH_2	25	15
COOH	25	83


Conclusions

1. Carboxyl group of glycine comes from carbon dioxide; methylene carbon comes from methanol
2. Hydroxymethyl group in serine comes from methanol; the other 2 carbons mimic the distribution seen in glycine
3. Serine arises by hydroxymethylation of glycine


Two possible routes for conversion of methanol plus CO₂ to cell material



NOTE: key difference is production of glycine by direct condensation (above) or by cleavage (below)



The serine cycle involves a cleavage reaction



What happens to the acetyl-CoA?

In icl⁺ bacteria: isocitrate lyase is involved in oxidation of acetyl-CoA to glyoxylate; in these bacteria ICL is also involved during growth on ethanol or acetate

In icl⁻ bacteria with no isocitrate lyase [eg *Methylobacterium extorquens*]

This route is not yet fully established.

It is also involved in metabolism of C2 compounds




Figure 3. The serine cycle as proposed by Peel, Large, Salem and Quayle⁹⁻¹¹. The enzymes are 1, serine transhydroxymethylase; 2, serine-glyoxylate aminotransferase; 3, hydroxypyruvate reductase; 4, glycerate kinase; 5, enolase; 6, PEP carboxylase; 7, malate dehydrogenase; 8, malate thiokinase; 9, malyl-CoA lyase. After the initial proposal much further enzymological and mutant evidence was subsequently accumulated to confirm this pathway³. Note that during biosynthesis of fatty acids and poly 3-hydroxybutyrate which use acetyl-CoA as their biosynthetic starting point, this pathway is sufficient for production of acetyl-CoA from formaldehyde plus carbon dioxide.




Figure 1 Kornberg's glyoxylate cycle⁴. This achieves the condensation of 2 molecules of acetyl-CoA to malate. The enzymes are: 1, citrate synthase; 2, aconitase; 3, isocitrate lyase; 4, malate synthase; 5, succinate dehydrogenase; 6, fumarase; 7, malate dehydrogenase. In effect this cycle can be summarized as $2 \text{ acetyl-CoA} \rightarrow \text{malate}$. A complete balance equation is given in Table 1 at the end of this review.





Figure 1 Kornberg's glyoxylate cycle⁴. This achieves the condensation of 2 molecules of acetyl-CoA to malate. The enzymes are: 1, citrate synthase; 2, aconitase; 3, isocitrate lyase; 4, malate synthase; 5, succinate dehydrogenase; 6, fumarase; 7, malate dehydrogenase. In effect this cycle can be summarized as $2 \text{ acetyl-CoA} \rightarrow \text{malate}$. A complete balance equation is given in Table 1 at the end of this review.



Confirmation of serine cycle

The proposed pathway fits the early labelled intermediates

The distribution of labelling in the intermediates fits the pathway


The 5 novel enzymes were purified and characterised

They were shown to be inducible on methanol


They were of sufficiently high specific activity to account for the growth rate on methanol

Mutants lacking them failed to grow on methanol; revertants had regained the enzyme

Later shown that key enzymes were coordinately regulated, implying the presence of an operon [Dunstan (Goodwin) & Anthony; Hanson & O'Connor (Lidstrom)]



Expression of the mxa operon



The genes: Nunn, Lidstrom, Amaratunga, Anderson, Anthony, Goodwin, Morris, O'Connor

[Top of carbon assimilation](#)
[Top of summaries](#)

For completion of the Serine cycle go to [Serine Cycle completion](#)

[Completion of the Serine /Ethylmalonyl-CoA Cycle](#) [PPT](#) [Top of summaries](#)

Solution of the Serine pathway in *Methylobacterium extorquens* (50 year project)

Situation in 1963 There must be a route for oxidation of acetylCoA to glyoxylate. An obvious route is to use the glyoxylate cycle but the key enzyme isocitrate is absent during growth on methanol (Large and Quayle 1963).

1970 Pat Dunstan (now Pat Goodwin). Showed that ICL is absent during growth on ethanol

Isolation of unusual mutants: A project to isolate MDH mutants. Penicillin enrichment procedure isolated mutants unable to grow on C1 or C2 but able to grow on succinate. These should be unable to oxidise methanol and ethanol. No mutants in assimilation pathways should be selected as these pathways are different.

Three types of mutant were isolated: MDH mutants. Cytochrome c mutants [unique, indicating the special role of cytochrome c in energy transduction from MDH]. A mutants with alteration in carbon assimilation pathways (PCT48) on C1 and C2 compounds showing there must be a common step in the pathway.




Figure 3. The serine cycle as proposed by Peel, Large, Salem and Quayle⁹⁻¹¹. The enzymes are 1, serine transhydroxymethylase; 2, serine-glyoxylate aminotransferase; 3, hydroxypyruvate reductase; 4, glycerate kinase; 5, enolase; 6, PEP carboxylase; 7, malate dehydrogenase; 8, malate thiokinase; 9, malyl-CoA lyase. After the initial proposal much further enzymological and mutant evidence was subsequently accumulated to confirm this pathway³. Note that during biosynthesis of fatty acids and poly 3-hydroxybutyrate which use acetyl-CoA as their biosynthetic starting point, this pathway is sufficient for production of acetyl-CoA from formaldehyde plus carbon dioxide.




Fig. 1. Kornberg's glyoxylate cycle⁴. This achieves the condensation of 2 molecules of acetyl-CoA to malate. The enzymes are: 1, citrate synthase; 2, aconitase; 3, isocitrate lyase; 4, malate synthase; 5, succinate dehydrogenase; 6, fumarase; 7, malate dehydrogenase. In effect this cycle can be summarized as 2 acetyl-CoA → malate.




Fig. 4. The serine cycle in methylotrophic bacteria having isocitrate lyase [ICL]³. The upper part of the Figure shows the serine cycle as shown on Fig 3. The lower part shows the oxidation of acetyl-CoA to glyoxylate by isocitrate lyase together with the non-decarboxylating enzymes of the TCA cycle.

Solution of the Serine pathway in *Methylobacterium extorquens* (50 year project)

Situation in 1963 There must be a route for oxidation of acetylCoA to glyoxylate. An obvious route is to use isocitrate lyase but this enzyme is absent during growth on methanol (Large and Quayle 1963).

1970 Pat Dunstan (now Pat Goodwin). Showed that ICL is absent during growth on ethanol

Isolation of unusual mutants: A project to isolate MDH mutants. Penicillin enrichment procedure isolated mutants unable to grow on C1 or C2 but able to grow on succinate. These should be unable to oxidise methanol and ethanol. No mutants in assimilation pathways should be selected as these pathways are different.

Three types of mutant were isolated: MDH mutants. Cytochrome c mutants [unique, indicating the special role of cytochrome c in energy transduction from MDH]. A mutants with alteration in carbon assimilation pathways (PCT48) on C1 and C2 compounds showing there must be a common step in the pathway.



Yuri Me, Pat Dunstan (now Goodwin) and Sasha Netrusov in Kiev 12 days after Chernobyl

The assimilation of ethanol in *M. extorquens* by study of ^{14}C -acetate assimilation

After growth on Methanol early label was in glycollate (reflects early glyoxylate label) & citrate


After growth on Ethanol early label was in glycine (reflects early glyoxylate label) & citrate

SO; there is an unknown common route for rapid metabolism of acetylCoA to glyoxylate during growth on C1 and C2 substrates.

In mutant 48 there was no rapid assimilation of acetate into glyoxylate (only citrate).

This same route was shown to operate on propanediol, 3-hydroxybutyrate and lactate

Problem: need to identify enzymes involved.



The shared pathway for methanol and ethanol assimilation




Fig. 5. Pathways for growth of *M. extorquens* on substrates metabolized by way of acetyl-CoA, based on the work of Pat Dunstan, John Bolbot and Ian Taylor^{12, 17-19, 21, 22}. NB: only the carbon balance is illustrated. Red indicates pathway on C₁ compounds; blue indicates pathway on C₂ and related compounds. In short-term labeling experiments glyoxylate would arise by equilibration with glyoxylate. The growth substrates include ethanol, acetate (a poor substrate), 3-hydroxybutyrate, malonate, propanediol, lactate and pyruvate. Propanediol and ethanol are oxidized by methanol dehydrogenase (MDH) whose electron acceptor is cytochrome c_L¹⁶; there is no growth of mutants lacking these proteins. For oxidation of propanediol by MDH an additional modifier protein is required to alter its substrate specificity²². Note that condensation of glyoxylate and acetyl-CoA to malate requires two enzymes: malyl-CoA lyase and malyl-CoA hydrolase.

Glyoxylate Regeneration Cycles

Mila Chistaserdova and Mary Lidstrom as a result of their work using mutants and some biochemistry produced many complex pathways, called Glyoxylate Regeneration cycles'

The solution was finally obtained in the lab of Georg Fuchs in Fribourg by very thorough enzymology and complex labelling techniques. Erb, Berg, Alber, Spanheimer, Ebenau-Jehle and Fuchs.

The EthylmalonylCoA pathway (EMC pathway)

This was done for acetate assimilation in *Rhodobacter sphaeroides* but was soon shown to be the common pathway also involved in methanol and ethanol assimilation in *M. extorquens*.


Most of the following slides are the Figures from my review:

How half a century of research was required to understand bacterial growth on C1 and C2 compounds: the story of the Serine Cycle and the Ethylmalonyl-CoA pathway. Science Progress 94, 109-138, 2011

Pathway based on Chistoserdova,
Lidstrom and colleagues

18 steps between acetyl-CoA and glyoxylate; how does ^{14}C -acetate glyoxylate so rapidly?

Biosynthesis



The Glyoxylate Regeneration Cycle Mila Chistaserdova and Mary Lidstrom



Mary Lidstrom (right)



Mila Chistaserdova (right)




Fig. 7. The glyoxylate regeneration cycle (GRC) for oxidation of acetyl-CoA in *M. extorquens* as proposed by Lidstrom, Chistoserdova and colleagues^{27, 31-33}. Their papers should be consulted for details of the extensive experimental work, mainly using mutants and radioactive substrates that led to this [necessarily] speculative proposal. The compounds in italics were later shown to be intermediates in the ethylmalonyl-CoA (EMC) pathway. The solid arrows merely indicate proposed reactions (or series of reactions); they do not necessarily indicate that such reactions are known reactions.




Fig. 9. Proposed pathway for acetyl-CoA assimilation by *Rhodobacter sphaeroides*.

This Figure is re-drawn from the 2006 paper by Alber, Spanheimer, Ebenau-Jehle and Fuchs⁴³. The gene *phaA* encodes β-ketothiolase; *phaB*, acetoacetyl-CoA reductase; *mch*, mesaconyl-CoA hydratase; *mcl1*, L-malyl-CoA/β-methylmalyl-CoA lyase; *pccAB*, propionyl-CoA carboxylase and *mcm* encodes (R)-methylmalonyl-CoA mutase. Although the enzymes catalyzing the conversion of the *C*₄ compound 3-hydroxybutyryl-CoA to the *C*₅ intermediate mesaconyl-CoA were not known at the time, it was suggested that this process probably involves a carboxylation step, as was subsequently demonstrated when the ethylmalonyl-CoA pathway was finally elucidated (Figs. 10-12).


Fig. 10. The 'missing' part of the ethylmalonyl-CoA (EMC) pathway. The conversion of crotonyl-CoA to to mesaconyl-CoA depends on three novel enzymes:


crotonyl-CoA carboxylase/reductase⁴⁴,

(2R)-ethylmalonyl-CoA mutase⁴⁶ and

(2)-methylsuccinyl-CoA dehydrogenase⁴⁷.

The two forms of ethylmalonyl-CoA are interconverted by ethylmalonyl-CoA/methylmalonyl-CoA epimerase.








Fig. 12. The ethylmalonyl-CoA (EMC) pathway for acetyl-CoA assimilation in *Rhodospirillum rubrum*. Note that there are two forms (R and S) of ethylmalonyl-CoA and two forms (R and S) of methylmalonyl-CoA (see Fig. 11) which are interconverted by the same epimerase.




Fig. 13. The serine/EMC cycle for assimilation of C₁ compounds by methylotrophs⁴⁴.

The ethylmalonyl-CoA (EMC) pathway for oxidation of acetyl-CoA to glyoxylate (lower half) (Fig. 12) is coupled to the serine cycle as shown on Fig. 3 (upper half). This is taken from the 2007 paper of Erb et al.⁴⁴ but for convenience only the carbon skeletons are shown. Dotted lines indicate that more than one reaction step is involved. Note that if acetyl-CoA is required as the biosynthetic precursor of membrane fatty acids or the storage compound poly 3-hydroxybutyrate then the EMC pathway is not required for oxidation of acetyl-CoA to glyoxylate.



Frieburg group: Georg Fuchs, Toby Erb and ? sorry



Celebrating
X, Georg Fuchs, Ivan Berg, Y, Z,
Toby

Sorry no picture of Birgit Alber




Fig. 14. **The serine/EMC cycle for assimilation of C1 compounds as it occurred during experiments described by Vorholt and colleagues⁵⁴** (re-drawn for ease of comparison with Figs. 3 and 13). This depiction of the pathway shows the succinyl-CoA, derived from propionyl-CoA, being 'recycled' to produce a third glyoxylate.



Julia Vorholt; confirmation of the Ethylmalonyl pathway (Zurich)

Two other cycles for acetate assimilaion




Fig. 6. The methylaspartate cycle²⁴. This pathway for oxidation of acetyl-CoA to glyoxylate in methylotrophs was proposed in 1984 by Shimizu, Ueda and Sato²³. Only the carbon skeletons have been included. The left hand side from mesaconyl-CoA to succinyl-CoA remains an essential part of the serine pathway as it is now understood. This cycle has recently been shown by Ivan Berg and colleagues to operate in haloarchaea for assimilation of C₂ compounds²⁴. In the complete methylaspartate cycle the glyoxylate condenses with a second molecule of acetyl-CoA to give malate, the overall carbon balance being the same as the glyoxylate cycle (Fig. 1).




Fig. 8. The citramalate cycle proposed in 1977 for oxidation of acetyl-CoA to glyoxylate in *Rhodospirillum rubrum* by Ivanovsky's group in Moscow (note: citramalate is α-methylmalate)^{39,40}. The pathway is completed by the condensation of the glyoxylate with a second acetyl-CoA to give malate.

[Top of Serine/EMC cycle](#)

[Top of summaries](#)

HOME

CONTACT ME: [Email](#)

THANKS for help to [Hugh Anthony](#) and [Carl Gore](#)

 [logo design](#)